



Contents lists available at ScienceDirect

Veterinary Parasitology

journal homepage: www.elsevier.com/locate/vetpar

Neosporosis in animals—The last five years

J.P. Dubey^{a,*}, G. Schares^b^a Animal Parasitic Diseases Laboratory, Animal and Natural Resources Institute, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD 20705, USA^b Institute of Epidemiology, Friedrich-Loeffler-Institut - Federal Research Institute for Animal Health, Seestrasse 55, D-16868 Wusterhausen, Germany

ARTICLE INFO

Keywords:

Neospora caninum
Biology
Diagnosis
Pathogenesis
Control

ABSTRACT

Neospora caninum is a protozoan parasite of animals. Until 1988, it was misdiagnosed as *Toxoplasma gondii*. Since its first recognition in 1984 in dogs and the description of a new genus and species *Neospora caninum* in 1988, neosporosis has emerged as a serious disease of cattle and dogs worldwide. Abortions and neonatal mortality are a major problem in livestock operations and neosporosis is a major cause of abortion in cattle. This review is focused on current status of neosporosis in animals based on papers published in the last five years. Worldwide seroprevalences are tabulated. Strategies for control and prevention are discussed.

© 2011 Elsevier B.V. All rights reserved.

Abbreviations: BIOVET, BIOVET-*Neospora caninum*, indirect ELISA, sonicate lysate of tachyzoites, BIOVET Laboratories, Canada; BioX, NcSRS2 sandwich ELISA, BioX, Belgium; CHEKIT, CHEKIT *Neospora*, indirect ELISA, detergent lysate of tachyzoites, IDEXX Laboratories, The Netherlands; CIVTEST, CIVTEST BOVIS NEOSPORA, indirect ELISA, sonicate lysate of tachyzoites, Laboratorios Hipra S.A., Spain; ELISA, enzyme linked immunosorbent assay; H, histology; IB, immunoblotting; IDEXX, IDEXX HerdChek *Neospora caninum* antibody, indirect ELISA, sonicate lysate of tachyzoites, IDEXX Laboratories, USA; ID-VET, ID SCREEN *Neospora caninum* indirect, indirect ELISA, ID-VET, France; IFAT, indirect fluorescent antibody test; IH, in house; IHC, immunohistochemistry; IH-ISCOM, detergent extracted tachyzoite antigen incorporated in immune stimulating complex particles; IH-Ncp43P, recombinant NcSRS2; IH-p38, native immune-affinity purified surface antigen NcSRS2; IH-NcSAG1, recombinant NcSAG1; IH-tNcSAG1, truncated recombinant NcSAG1; IH-NcSRS2, recombinant NcSRS2; IH-tNcSRS2, truncated recombinant NcSRS2; MASTAZYME, MASTAZYME NEOSPORA, indirect ELISA, formaldehyde-fixed whole tachyzoites, MAST GROUP UK; NS, not stated; NAT, *Neospora* agglutination test; NhSAG1, recombinant NhSAG1; PCR, polymerase chain reaction; Pourquoi, Institut Pourquoi, Montelancier, France; SVANOVA, commercialized ISCOM ELISA, SVANOVA Biotech AB, Sweden; VMRD, *Neospora caninum* cELISA Competitive ELISA GP65 surface antigen of tachyzoites VMRD, USA; WH, whole tachyzoite extract; WT-IHCA, kinetic ELISA-California, USA.

* Corresponding author. Tel.: +1 301 504 8128; fax: +1 301 504 9222.

E-mail address: jitender.dubey@ars.usda.gov (J.P. Dubey).¹ J.P. Dubey is employee of the US Department of Agriculture, and the work is part of his official duty.

1. Introduction

Neospora caninum is morphologically similar to *Toxoplasma gondii* but these parasite species are biologically different. Neosporosis, is primarily a disease of cattle and dogs and is not considered zoonotic whereas toxoplasmosis is a serious disease of humans, sheep, and many other warm-blooded animals. Although much has been published on the biology of *N. caninum* during the last 23 years since its discovery in 1988 (Dubey et al., 2002), neosporosis continues to be a major problem in cattle. In the present review we summarize information on neosporosis in animals for the past 5 years.

2. General biology

2.1. Host range

Past (Dubey et al., 2007a) and recent surveys (Tables 1–9) indicate that a wide range of domestic and wild animals have been exposed to *N. caninum*. However, viable *Neospora* has been isolated from only a few hosts (cattle, sheep, water buffalo, dog, horse, bison, white-tailed deer); recent reports on isolations are summarized in Table 1.

Table 1
Recent viable isolates of *N. caninum*.

Hosts	Country	Tissue/origin	No. of isolates (designation)	Reference
Cattle (<i>Bos taurus</i>)	Brazil	Brain of an asymptomatic 4-month old calf	1 (Nc-Goiás 1)	García-Melo et al. (2009)
	Israel	Fetal brain	2 (NcIs491, NcIs580)	Fish et al. (2007)
	Poland	Asymptomatic calf brain	NC-PolB1	Goździk and Cabaj (2007)
	Slovakia	Adult cow	Nc-SKB1	Reiterová et al. (2011)
	Spain	Brain of asymptomatic calves	9 (Nc-Spain 6,7,8,9,10, Nc-Spain 2H,3H, 4H, 5H)	Regidor-Cerrillo et al. (2008)
European bison (<i>Bison bonasus bonasus</i>)	Poland	Brain of asymptomatic calf	1 (Nc-Spain 1H)	Rojo-Montejo et al. (2009b)
		Blood	2 (NC-PolBb1 and 2)	Bień et al. (2010)
Dog (<i>Canis familiaris</i>)	Germany	Feces, bioassay in KO mice and cell culture	3 (NC-GER7, 8, 9)	Basso et al. (2009b)
	Portugal	Feces, bioassay in KO mice and cell culture	1 (NC-P1)	Basso et al. (2009a)

Table 2
Small mammals and birds as natural hosts for *N. caninum*.

Host	Country or region	Remarks	Reference
Avians			
Chicken (<i>Gallus domesticus</i>)	Americas ^a	Antibodies (IFAT, 1:25 or higher) detected in 39.5% of 1324 free range chickens	Martins et al. (in press)
	Brazil	DNA detected in 6 of 10 seropositive chickens Antibodies (IFAT, 1:50) detected in 23.5% of 200 free-range, and 1.5% of 200 indoors chickens	Costa et al. (2008)
Sparrow (<i>Passer domesticus</i>)	Brazil	DNA detected in 3 (7.5%) of 40 sparrows	Gondim et al. (2010)
Rodents			
Field mouse (<i>Apodemus sylvaticus</i>)	Italy	DNA detected in brain of 1, and muscle of 1 of 55 field mice from North West	Ferroglio et al. (2007a)
Rat (<i>Rattus norvegicus</i>)	Italy	DNA detected in brains of 2, kidneys of 4, and muscles of 10 of 103 rats from North West	
House mouse (<i>Mus musculus</i>)	Australia	DNA detected in 26.9% of 104 mice from Sydney and in 8 mice in extra neural tissues	Barratt et al. (2008)
	Italy	DNA detected in brain of 2, kidney of 1, and muscle of 8 of 75 mice from North West	Ferroglio et al. (2007a)
Capybara (<i>Hydrochaeris hydrochaeris</i>)	Brazil	DNA detected in 3 (11.5%) of 26; in lymph node of 2 and liver of 1 capybara	Truppel et al. (2010)
		Antibodies (IFAT, 1:50) detected in 9.4% of 213 feral capybaras	Yai et al. (2008)
		Antibodies (IFAT, 1:50) detected in 39% of 63 feral capybaras	Valadas et al. (2010b)
Vole (<i>Microtus arvalis</i>)	Austria	DNA detected in brain of 4 (1.5%) of 264 voles	Fuehrer et al. (2010)
Water vole (<i>Arvicola terrestris</i>)	Austria	DNA detected in brain of 2 (2.3%) of 86 water voles	Fuehrer et al. (2010)
Lagomorphs			
Rabbit (<i>Oryctolagus cuniculus</i>)	UK	DNA detected in brain of 6 (10.5%) of 57 wild rabbits from Yorkshire	Hughes et al. (2008)
	Egypt	Antibodies (IH-tNcSAG1) found in 1.8% of 54 farm rabbits	Ibrahim et al. (2009)
	Austria	Antibodies (VMRD) found in 37% of 383 hares	Bártová et al. (2010b)
Hare (<i>Lepus europaeus</i>)	Czech Republic	Antibodies (VMRD) found in 39.7% of 333 hares	

^a Antibodies to *N. caninum* were detected in 18.5% of 97 chickens from Mexico, 7.2% of 97 chickens from USA, 39.5% of 144 chickens from Costa Rica, 71.5% of 102 chickens from Grenada, 44% of 50 chickens from Guatemala, 83.6% of chickens from Nicaragua, 58.1% of 55 chickens from Argentina, 34.3% of 358 chickens from Brazil, 62.3% of 85 chickens from Chile, 11.2% of 62 chickens from Colombia, 38.7% of 80 chickens from Guyana, 18% of 50 chickens from Peru, and 21.7% of 46 chickens from Venezuela (Martins et al., in press).

Unlike *T. gondii*, it is difficult to isolate viable *N. caninum*. Additionally, some isolates were obtained as oocysts by feeding naturally infected tissues to dogs and these isolates were not cryopreserved for future studies. For example, to our knowledge none of the isolates from water buffalo from Brazil were grown in cell culture. Additionally, not all isolates of *N. caninum* could be cultivated in cell culture

(Vianna et al., 2005). We emphasize that finding DNA is not comparable with finding viable *N. caninum*. It is notable that *N. caninum* DNA was demonstrated frequently in tissues of asymptomatic rodents but viable parasite has not been isolated (Table 2).

Humans are not regarded as intermediate host of *N. caninum* (McCann et al., 2008).

Table 3
Prevalence of *N. caninum*-like oocysts in feces of dogs.

Country	No. of dogs	Type	No. positive	Microscopic	NC-PCR	Bioassay	Reference
Costa Rica	34	Dairy farms	3	0	3	0	Palavicini et al. (2007)
Iran	174	89 farm, 85 house-hold	4	4	2	Not done	Razmi (2009)
Italy	230	Farms	0	0	Not done	Not done	Paradies et al. (2007)
Spain	285	Farms	1	1	Not done	Not done	Regidor-Cerrillo et al. (2010a)

Table 4Prevalence of *N. caninum* antibodies in dogs.

Country	Type	No. tested	% positive	Test	Titer/Supplier	Reference
Algeria		100	21.0	ELISA	BioX	Ghalmi et al. (2009b)
	Pound	261	22.5	IFAT	NS	Ghalmi et al. (2009a)
	Police	85	6.6			
	Breeder	184	12			
	Farm	80	44.4			
Brazil						
Bahia, Salvador	Urban	49	32.7	Wb, ELISA	VMRD	Jesus et al. (2007)
Goiás	Urban	197	32.9	IFAT	1:50	Boaventura et al. (2008)
Mato Grosso	Clinics	60	45.0	IFAT	1:50	Benetti et al. (2008)
	Dairy farms	37	67.6	IFAT	1:200	Benetti et al. (2009)
Minas Gerais	Clinics	228	3.1	IFAT	1:50	Guimarães et al. (2009)
Pará	Rural	72	11.1	IFAT	1:50	Valadas et al. (2010a)
	Urban-stray	57	14			
Paraná state	Urban	181	12.7	IFAT		Fridlund-Plugge et al. (2008)
	Periurban	178	15.7		1:50	
	Rural	197	25.3			
Pernambuco						
Paulista	Domiciled	289	26.0	IFAT	1:50	Figueredo et al. (2008)
Amaraji	Domiciled	168	26.2			
Garanhuns	Domiciled	168	34.5			
Piauí	Urban	530	30.2	IFAT	1:50	Lopes et al. (2011)
Rio Grande do Sul	Rural	230	20.4	IFAT	1:50	Cunha Filho et al. (2008)
	Urban	109	5.5	IFAT	1:50	
São Paulo	Urban	108	15.7	IFAT	1:50	Bresciani et al. (2007) ^a
	Beef farms	963	25.4	IFAT	1:50	de Moraes et al. (2008)
Canada						
Northwest Territories	Clinics	108	3.7	IFAT	1:25	Salb et al. (2008)
Costa Rica	Farm	31	48.4	cELISA	VMRD	Palavicini et al. (2007)
Grenada, West Indies		107	2	IFAT	1:100	Dubey et al. (2008b)
India	Rural	126	21.4	ELISA	VMRD	Sharma et al. (2008)
	urban	58	6.9			
Iran	Farm	50	28	IFAT	1:50	Haddadzadeh et al. (2007)
	Urban	50	11.3	IFAT	1:50	
	Clinics	233	10.3	ELISA	P38	Hosseininejad et al. (2010)
				IFAT	1:50	
Urmia	Stray	135	27	IFAT	1:50	Yakhchali et al. (2010)
Italy	Urban	188	20.2	36.4 NAT	1:40	Ferroglio et al. (2007b)
	Rural	302				
	Kennel	144	14.6	ELISA	MASTAZYME	Paradies et al. (2007) ^a
	Farm	162	26.5	ELISA	MASTAZYME	Paradies et al. (2007)
Japan	Clinics	1206	10.4	ELISA	IH-tNcSAG1	Kubota et al. (2008)
Mexico						
Aguascalientes	Urban	116	20	ELISA	IDEXX	Cruz-Vázquez et al. (2008)
	Dairy farms	152	40.7			
Durango City	Pound	101	2	IFAT	1:25	Dubey et al. (2007b)
Peru	Farm	122	14.8	IFAT	1:50	Vega et al. (2010)
Poland	Clinics	257	21.7	ELISA		Goździk et al. (2011)
	Clinics	110	16.3	IFAT	1:50	Płoneczka and Mazurkiewicz (2008)
Senegal, West Africa		196	17.9	ELISA	VMRD	Kamga-Waladjo et al. (2010)
Spain						
Andalusia	Feral	28	17	ELISA	VMRD	Millán et al. (2009a)
Galicia	Farm	141	47.5	IFAT	1:50	Regidor-Cerrillo et al. (2010a,b) ^a
	Stray	134	39.5	IFAT	1:50	Regidor-Cerrillo et al. (2010a,b)
Majorca island	Kennel	44	0	ELISA	VMRD	Cabezón et al. (2010)
Several areas	House hold	102	2.9	IFAT	1:50	Collantes-Fernández et al. (2008) ^a
	Stray	94	24.5			
	Hunting	100	23			
	Farm	100	51			
Turkey						
Kirikkale	Stray	121	28.9	IFAT	1:16	Yildiz et al. (2009) ^a

NS, not stated.

^a Risk factors.

2.2. Definitive hosts and transmission by oocysts

Experimental studies showed that the domestic dog and the Australian dingo (both *Canis domesticus*), and

the coyote (*Canis latrans*) are definitive hosts for *N. caninum* (McAllister et al., 1998; Gondim et al., 2004; King et al., 2010). Of these, viable oocysts have been demonstrated only in feces of naturally-infected dogs (Basso

Table 5
Serologic prevalence of *N. caninum* antibodies in cattle.

Country	Region	No. of animal (relevant details)	% Positive	Assay	Cut-off titer or test	Reference
Algeria		102	3.9	ELISA	BioX	Ghalimi et al. (2009b)
Argentina		4190 no abortion	14.2	IFAT	1:200	Moore et al. (2009)
		1042 abortion	25.7	IFAT	1:200	Moore et al. (2008)
Brazil	Mato Grosso do Sul	173 dairy	80.9	IFAT	1:25	Moré et al. (2009)
		90 beef	73.0	IFAT	1:25	Moré et al. (2008a)
		1098 beef	62.5	IFAT	1:50	Andreotti et al. (2010) ^a
	Minas Gerais	2448	14.9	IFAT	1:50	Oshiro et al. (2007) ^a
		932 dairy	53.5	IFAT	1:200	Benetti et al. (2009)
		559 dairy	91.2	IFAT	1:200	Guedes et al. (2008)
		575 cows slaughter	97.2	IFAT	1:200	
	Pará	503 fetuses	12.7	IFAT	1:25	
		120 beef	19.2	IFAT	1:100	Minervino et al. (2008)
	Paraná	40 dairy	17.5	IFAT	1:100	Minervino et al. (2008)
		159 beef (<i>Bos indicus</i>)	15.1	ELISA	IDEXX	Marques et al. (2011)
Pernambuco		469	31.7	IFAT	1:200	Silva et al. (2008) ^a
Rio de Janeiro		563 dairy	23.3	ELISA	IDEXX	Munhoz et al. (2009) ^a
Egypt		93	20.4	ELISA	IH-tNcSAG1	Ibrahim et al. (2009)
Estonia		320 bulk milk	16.0	ELISA	SVANOVA	Lassen et al. (2008)
Germany		1950 bulk milk	1.0	ELISA	IH-p38	Schares et al. (2009)
Greece		1573 milk	15.2	ELISA	IH-p38	Sotiraki et al. (2008)
Iran		285	12.6	ELISA	SVANOVA	Fard et al. (2008)
		237	32.0	ELISA	IDEXX	Youssefi et al. (2009)
Mexico	Nuevo Leon	813	11.6	ELISA	NS	Segura-Correa et al. (2010)
	Veracruz state	863	26.0	ELISA		Romero-Salas et al. (2010)
	4 provinces	596	11.6	ELISA	IDEXX	Garcia-Vazquez et al. (2009)
Norway		1657 herds bulk milk	0.7	ELISA	SVANOVA	Klevar et al. (2010)
Pakistan	Punjab province	240	43.8	cELISA	VMRD	Shabbir et al. (in press)
Peru	Department Junín	347	12.4	IFAT		Puray et al. (2006)
People's Republic of China		300	20.3	ELISA	IH-NcSRS2	Liu et al. (2007)
		540 dairy	13.3	ELISA	IDEXX	Wang et al. (2010)
	Beijing	212	43.4	ELISA	IH-tNcSRS2	Yao et al. (2009)
	Tianjin	601	5.7	ELISA	IH-tNcSRS2	Yao et al. (2009)
	South	370 dairy	18.9	ELISA	IDEXX	Xia et al. (2011)
Philippines		96	16.7	ELISA	IH	Konnai et al. (2008)
Romania	Cluj, Satu-Mare, Mureş, Sibiu, Alba	193	55.9	ELISA	IDEXX	Gavrea and Cozma (2010)
Spain	Galicia	37,090 dairy	22.5	ELISA	IDEXX	Eiras et al. (2011) ^a
		20,206 beef	25.6			
		2292 mixed	25.4			
		5196	15.7	IFAT	1:50	González-Warleta et al. (2008)
		178	7.3	c-ELISA	VMRD	Panadero et al. (2010)
Slovakia		716	20.1	ELISA	Pourquier	Reiterová et al. (2009)
Sweden		2754	2.8	ELISA	IH-ISCOM	Loobuyck et al. (2009)
Thailand	Khon Kean	424	8.0	ELISA	IH-ISCOM	Chanlun et al. (2007)
Turkey		25 aborted	60.0	ELISA	VMRD	Kul et al. (2009)
		40 heifers	40.0			
		6 calves	33.3			
		89 repeat breeder	13.4	ELISA	VMRD	Simsek et al. (2008)
		94 healthy	3.1			
		234 aborted	6.8	ELISA	VMRD	Yildiz et al. (2009)
		323 no abortion	10.7			
		15736	12.9	ELISA	IDEXX	Woodbine et al. (2008)
United Kingdom		460 dairy heifers	7.2	ELISA	MASTAZYME	Brickell et al. (2010)
		900 beef	16.7	Kinetic ELISA	WT-IHCA	Hoar et al. (2007)
USA		254 (milk)	30.0	ELISA	SVANOVA	Geurden et al. (2008)
Vietnam		215 dairy	41.0	ELISA	IH-ISCOM	Duong et al. (2008)

NS, not stated.

^a Risk factors.

et al., 2009a,b), and recently in naturally-infected gray wolf (*Canis lupus*) (Dubey et al., in press). In foxes no viable oocysts have been observed yet, and in a survey in Germany no *N. caninum* oocyst were detected in fox feces (Constantin et al., 2011).

Oocysts are the key in the epidemiology of neosporosis. They are environmentally resistant like the oocysts of other coccidians (Uzeda et al., 2007; Neto et al., 2011).

N. caninum oocysts have been identified in only a few dogs worldwide. The number of oocysts shed by dogs is usually low. No or only a few oocysts were seen in recent surveys of canine feces (Table 3). Because *N. caninum* oocysts structurally resemble another coccidian in dog feces, *Hammondia heydorni*, it is epidemiologically important to properly identify *N. caninum* oocysts (Soares et al., 2011). Recently additional molecular tools for the differ-

Table 6

Diagnosis of *N. caninum*-associated abortion in dairy cattle from selected studies based on fetal examination.

Country	No. of fetuses examined	% Infected	Reference
Argentina	666	9.9 ^a	Moore et al. (2008)
Brazil	258	34.0 ^b	Pescador et al. (2007)
Germany	232	10.0 ^c	Sörgel et al. (2009)
Iran	100	13.0 ^d	Razmi et al. (2007)
	151	14.5 ^e	Razmi et al. (2010)
	12	100 ⁱ	Salehi et al. (2009)
Japan	15	27.0 ^h	Ghanem et al. (2009)
People's Republic of China	16	4.0 ^h	Zhang et al. (2007)
	26	57.7 ^j	Yao et al. (2009)
Romania	9	33.0 ^j	Suteu et al. (2010)
Switzerland	223	16.1 ^f	Reitt et al. (2007)
	58	40.0 ^g	Tschuor et al. (2010)

^a 66 Positive by at least 1 method. Lesions in 70; 49 of these positive by IHC; 31 of 55 fetal serology positive. DNA detected in frozen tissues in 17 of 17 fetuses, and in 17 of 53 in fetuses from paraffin-embedded tissues. The positive PCR tissues were: liver 54.4%, CNS 24.2%, placenta 33.3%, and heart 10.5%.

^b Lesions in 89 fetuses, IHC positive in 55 fetuses, lungs and brain most useful for histology.

^c IHC (in 16), PCR (in 16–19), ELISA (in 14).

^d 13 Fetuses PCR positive, lesions 12 fetuses, 3 IHC positive.

^e 18 (11.9%) PCR positive, 6 of 52 IHC positive.

^f Histologic lesions 71.4%, IHC 8.6%, nested PCR 6.7%, nested PCR-PLUS 20.9%.

^g Maternal antibodies, histological examination of brain and heart of the fetus, PCR of fetal brains.

^h PCR, in one case tissue cyst observed.

ⁱ 12 of 12 nested PCR positive; 3 of 12 histologically positive in brain.

^j PCR.

entiation of *N. caninum* and *H. heydorni* were reported and applied to rodent tissues (Barratt et al., 2008).

How dogs become infected with *N. caninum* in nature is not fully understood. Historically, vertical transmission of neosporosis was first recognized in dogs. *N. caninum* is considered to be transmitted from the dam to the neonates during terminal stages of gestation or post-natally via milk. Unlike cattle, vertical transmission of *N. caninum* in dogs is considered highly variable and not likely to persist in the absence of horizontal infection.

Fecal transmission of *N. caninum* in dogs appears to be less important than carnivorous. Bandini et al. (2011) fed 4 dogs with 1000, 5000 or 10,000 *N. caninum* oocysts; none of the 4 dogs shed *N. caninum* – like oocysts in their feces during the observation period of 30 days. However, the 2 dogs fed with 10,000 oocysts seroconverted but the 2 dogs fed with 1000 or 5000 oocysts did not. Neither parasite DNA nor the parasite stages were demonstrable in tissues of the seropositive dogs euthanized 6 months after feeding oocysts. These findings suggest fecal transmission may not be an important mode of transmission of the parasite for the definitive host but results need confirmation.

Age-related prevalence data indicate that most dogs become infected after birth; higher prevalences have been documented in older versus younger dogs. The ingestion of infected tissues is the most likely source of infection for carnivores. Theoretically, tissues of any animal containing tissue cysts can be a source of infection for dogs. Tissues of infected prey of dogs may represent a logi-

cal source of infection; but viable parasite has not been isolated from potential dog prey as e.g. birds, rodents or lagomorphs (Table 2). Experimentally, chickens older than one week inoculated with tachyzoites intraperitoneally developed transient infection. Parasites or antibodies were not demonstrable 60 days p.i. However, inoculation of chicken embryonated eggs produced patent infection and the infected chorioallantoic membranes of these eggs induced oocyst shedding when fed to a dog (Furuta et al., 2007). The susceptibility of chicken eggs for *N. caninum* infection was recently confirmed by Mansourian et al. (2009) using broiler chicken embryonated eggs. Pigeons (*Columba livia*) were also successfully infected with *N. caninum* tachyzoites and are also putative natural reservoirs for *N. caninum* (Mineo et al., 2009). Concerning natural infections in wild birds, in a recent work, *N. caninum* DNA has been observed in brain from magpies (*Pica pica*) and common buzzard (*Buteo buteo*), including these species as natural intermediate host for *N. caninum* (Darwich et al., submitted for publication).

Higher seroprevalences of *N. caninum* antibodies in rural dogs versus city dogs (Table 4) is probably related to availability of prey or infected animals for carnivorous. Improper disposal of dead infected cattle on the farm or wild life can be a source of infection. Both neural and extra-neural tissues can be a source of infection for dogs. Dogs fed masseter muscle, heart, liver, and brains of naturally-infected cattle or buffaloes shed oocysts (Bandini et al., 2011; Cavalcante et al., 2011). Until now *N. caninum* tissue cysts have been demonstrated only in neural and muscular tissues. Whether tissues containing only tachyzoites (acutely infected animals) can be orally infectious to animals is not known. Dogs fed tissues from infected bovine neonates did not shed oocyst (Cedillo et al., 2008), maybe because *N. caninum* often dies together with the host tissue.

2.3. Strain variation and virulence

It is now well established that *N. caninum* can cause serious illness in cattle and dogs, and occasionally in other animals. Infections in many hosts are common but clinical disease is rare. Clinical disease maybe associated with the strain of *N. caninum*. Isolates of *N. caninum* from various hosts are genetically similar although many strains have their own molecular signature as determined by multilocus microsatellite analysis (Regidor-Cerrillo et al., 2006; Al-Qassab et al., 2009a, 2010; Basso et al., 2009b, 2010). The molecular characteristics of a strain could be useful for epidemiological studies. Exogenous point source outbreaks of bovine neosporosis in Germany were associated with a common source of infection (Basso et al., 2010). In another investigation, there was clustering of *N. caninum* isolates according to geographical origin of aborted bovine fetuses in Spain (Pedraza-Díaz et al., 2009). Little is known of the strain variation with respect to their virulence. In limited studies some *N. caninum* strains were more virulent to mice than others and showed also differences during *in vitro* cultivation (García-Melo et al., 2009, 2010; Rojo-Montejo et al., 2009a,b; Regidor-Cerrillo et al., 2010b, 2011). It is not yet known, whether virulence in mice could reflect the effect an *N. caninum* infection in other host.

Table 7
Prevalence of antibodies to *N. caninum* in sheep and goats.

Host	Country	No. examined	% Positive	Assay	Cut-off titer or test	Reference
Sheep (<i>Ovis ovis</i>)	Australia	232	2.2	ELISA	VMRD	Bishop et al. (2010)
	Brazil					
	Alagoas	343	9.6	IFAT	1:50	Faria et al. (2010) ^a
	Campo Grande	441	30.8	IFAT	1:50	Andreotti et al. (2009)
			32.0	ELISA	IH-NcSRS2	
	Federal District	1028	8.8	IFAT	1:50	Ueno et al. (2009)
	Minas Gerais State	155	47.1	IFAT	1:64	Rossi et al. (2011)
		334	8.1	IFAT	1:50	Salaberry et al. (2010)
	Rio Grande do Norte	409	1.8	IFAT	1:50	Soares et al. (2009)
	Roraina	141	29.0	IFAT	1:50	Aguiar et al. (2004)
	São Paulo	382	12.8	IFAT	1:25	Langoni et al. (2011)
	Czech Republic	547	12.0	ELISA	VMRD	Bártová et al. (2009)
	Jordan ^a					
	North	339	63.0	ELISA	BioX	Abo-Shehadeh and Abu-Halaweh (2010) ^a
	South	320	4.3	ELISA	CHEKIT	Al-Majali et al. (2008) ^a
	New Zealand	137 (aborted)	37.0	IFAT	1:100	Howe et al. (2008)
		640	0.6	ELISA, IFAT	1:50	Reichel et al. (2008)
	Philippines	38	26.3	IB,ELISA	IH	Konnai et al. (2008)
	Slovakia	382	3.7	ELISA	ID-VET	Špilovská et al. (2009), Špilovská and Reiterová (2008)
	Goat (<i>Capra hircus</i>)	Spain				
Galicia		177	57.0	ELISA	VMRD	Panadero et al. (2010)
Argentina		1594	6.6	IFAT	1:50	Moore et al. (2007)
Brazil						
Paraíba state		306	3.3	IFAT	1:50	Faria et al. (2007)
Rio Grande do Norte		381	1.0	IFAT	1:50	Lima et al. (2008)
São Paulo		923	19.7	NAT	1:25	Modolo et al. (2008)
Jordan ^a						
North		302	2.0	ELISA	BioX	Abo-Shehadeh and Abu-Halaweh (2010) ^a
South		300	5.7	ELISA	CHEKIT	Al-Majali et al. (2008) ^a
People's Republic of China		207 cashmere	7.7	ELISA	IH-NcSAG1	Lu et al. (2007)
Philippines		89	23.6	ELISA	IH	Konnai et al. (2008)
Poland		1060	0.4	ELISA	VMRD	Czopowicz et al. (2011)
Slovakia	18	16.6	ELISA	ID-VET	Špilovská and Reiterová (2008)	

Abortion or fetal infections have been induced in cattle using a variety of isolates in different laboratories but a meaningful comparison in pregnant cattle will be economically prohibitive to do (Dubey et al., 2007a). In a recent study, cows inoculated with a Spanish bovine isolate did not damage the fetus even though cows were inoculated at an early (70 days) stage of gestation age (Rojo-Montejo et al., 2009a). On the other hand, in beef cattle experimentally infected at 110 days of gestation with a beef strain, fetal death occurred in one fetus (Almería et al., 2010).

Studies aiming to examine virulence of *N. caninum* in cattle are difficult to interpret. Because experimentally infected dogs usually shed only few oocysts, this stage is often not available. Therefore, often tachyzoites are used for experimental infection of cattle although oocysts are the most likely source of postnatal infection in cattle and the outcome of disease might differ depending of the parasite stage by which an animal becomes infected. Studies using cell culture derived tachyzoites may further be complicated because *N. caninum* isolates maintained *in vitro* for long time may be altered regarding their virulence and other biological characteristics. Experimental induction of abortion may vary with the strain, the passage number of *N. caninum* or the route of inoculation; e.g. none of the 19 heifers inoculated intraconjunctively with 10⁸ NC-1 tachyzoites delivered an infected calf (de Yaniz et al., 2007).

3. Neosporosis in cattle

3.1. Serologic prevalence

Serologic prevalences of *N. caninum* summarized in Table 5 indicate that there are considerable differences among countries, within countries, between regions, and between beef and dairy cattle. However, caution should be used in evaluating these results because of differences in serologic techniques, study design, and sample size used (Dubey et al., 2007a). Most of these surveys were made using individual cattle sera. Bulk milk serology is an economical way of estimating *N. caninum* prevalence on a herd basis (Wapenaar et al., 2007b; Frössling et al., 2008; Schares et al., 2009), but this method is not as accurate as the detection of antibodies in the serum.

There are indications that the *N. caninum* seroprevalence differs according to the cattle breed (Armengol et al., 2007; Duong et al., 2008; Munhoz et al., 2009). However, some of these results might have been caused by differences in the production system used for the different breeds and not by differences in the breed-related susceptibility to infection. In a study with large sample size, seroprevalences in beef cattle (25.6% of 20,206) and dairy cattle (22.5% of 37,090) were similar (Eiras et al., 2011). Whether improved breeds of dairy cattle are more susceptible to *N. caninum* infection than the zebu cattle or cross breeds needs further investigation (Munhoz et al., 2009).

Table 8Prevalence of antibodies to *N. caninum* in miscellaneous domestic animals.

Host	Country	No. examined	% Positive	Assay	Cut-off titer or test	Reference
Domestic cat (<i>Felis catus</i>)	Spain	20	15.0	ELISA	VMRD	Millán et al. (2009a)
	Hungary	330	0.6	IFAT	1:40	Hornok et al. (2008)
Camel (<i>Camelus dromedarius</i>)	United Arab Emirates	1119	13.7	ELISA	VMRD	Wernery et al. (2008)
South american camelids	Argentina	308	4.6	IFAT	1:25	Moré et al. (2008b)
Llama (<i>Lama glama</i>)	Peru	175	2.9	IFAT	1:100	Casas et al. (2006)
Pig (<i>Sus scrofa</i>)	Senegal, West Africa	60 wild	58.3	ELISA	VMRD	Kamga-Waladjo et al. (2009)
	Brazil	130	3.1	IFAT	1:50	Azevedo et al. (2010)
Yak (<i>Bos grunniens</i>)	People's Republic of China	946	2.2	ELISA	IDEXX(Herd Check)	Liu et al. (2008)
Bali cattle (<i>Bos javanicus</i>)	Indonesia	438	5.5	ELISA	IH-p38	Damriyasa et al. (2010)
Gaur (<i>Bos gaurus f. frontalis</i>)	India	159	10.0	ELISA	VMRD	Rajkhowa et al. (2008)
Water buffalo (<i>Bubalus bubalis</i>)	Argentina	449	64.0	IFAT	1:100	Campero et al. (2007)
	Iran	181	37	ELISA	IDEXX	Hajikolaie et al. (2007)
	Pakistan	300	54.7	ELISA	VMRD	Nasir et al. (2011)
	Philippines	105	3.8	ELISA	IH	Konnai et al. (2008)

In this respect there are very few reports of *N. caninum* infection in indigenous breeds of cattle from Asia. In a survey from Pakistan, *N. caninum* seropositivity was lower in European dairy breeds than in cross bred and non-descript cattle (Shabbir et al., in press). Rate of abortion and immune responses after *N. caninum* infection might also be affected by the breed of cattle (Armengol et al., 2007; Almería et al., 2009b; Romero-Salas et al., 2010; Santolaria et al., 2011).

Little information is available on the association of genetic traits and *N. caninum* infection. In a retrospective study of Holstein cattle in Canada, Schwab et al. (2009) found no association between *N. caninum* seropositivity and allele frequency distribution of DRB3 or DQA genes. However, independent of the serological status the alleles DRB3*1001 and DRB3*2703 were associated with resistance and susceptibility of pregnancy loss.

One of the short coming of serological surveys is the validity of the techniques used. As of yet, neither intact *N. caninum* nor parasite DNA were demonstrable in asymptomatic adult cows (Dubey et al., 2007a; Wapenaar et al., 2007a; Moré et al., 2008a; Santos et al., 2010). However, *N. caninum* could be isolated from the brains of asymptomatic naturally infected calves (Fish et al., 2007; Goździk and Cabaj, 2007; Regidor-Cerrillo et al., 2008; García-Melo et al., 2009; Rojo-Montejo et al., 2009b).

3.2. Transmission and epidemiology

Vertical transmission from the dam to the fetus, and post natal ingestion of oocysts are the only demonstrated modes of transmission in cattle. *N. caninum* is one of the most efficiently transplacentally transmitted parasites among all known microbes in cattle. Transplacental transmission can occur in postnatally acquired infections by ingestion of oocysts (exogenous) or reactivation of infection in a chronically infected cow (endogenous) and the rate of transmission may differ in these two scenarios (Williams et al., 2009).

The rate of vertical transmission may vary among herds. In selected Dutch dairy herds vertical transmission rates were estimated to be 61.8% (Bartels et al., 2007) in one study, and 58% in another one (Dijkstra et al., 2008). Vertical transmission rate was 37.1% in a dairy herd in Argentina; cows with high titers had more infected calves

than cows with low titers (Moré et al., 2009). Serological examination of precolostral serum is a convenient way of measuring congenital infection. In the absence of active infection, transcolostrally-acquired antibodies decay with time depending on the titer in the colostrums and the detection of colostral antibodies in calves is also influenced by the sensitivity of serological test. In one study, passively acquired *N. caninum* antibodies persisted for five weeks (Cardoso et al., 2008).

The ingestion of sporulated *N. caninum* oocysts from the environment is the only demonstrated natural mode of infection in cattle after birth (McCann et al., 2007). To date cow to cow transmission of *N. caninum* has not been observed. In a very large sample size involving 37,090 dairy and 20,206 beef cattle in Spain, seroprevalence increased with age (Eiras et al., 2011). Seroprevalences in dairy cattle were: 10.4% in 12–24 months old, 14.1% in 25–36 months old, and 24.6% in >36 months old; similar increase was seen in beef cattle (12.9%, 15.3%, and 31.8% in the 3 age groups, respectively). These data indicate significant post-natal transmission in the area under examination.

Other modes of transmission suggested are via milk, and semen. Venereal transmission is possible, but unlikely since under experimental conditions large numbers of tachyzoites were necessary for infection (Serrano-Martínez et al., 2007b; Ferre et al., 2008). Dams naturally bred with experimentally infected bulls failed to seroconvert (Osoro et al., 2009).

3.3. Clinical neosporosis

N. caninum is a major cause of abortion in both dairy and beef cattle, and worldwide prevalences were summarized (Dubey, 2003a; Dubey et al., 2007a). Cows of any age may abort from three months gestation to term with most abortions occurring at five to six month gestation. Fetuses may die in utero, be resorbed, mummified, autolyzed, still-born, born alive with clinical signs, or born clinically normal but persistently infected. Recently, *N. caninum* was demonstrated in tissues of 4 of 15 mummified fetuses (Ghanem et al., 2009); these fetuses tested negative for SLC35A3 gene that causes complex vertebral malformation in cattle, suggesting *N. caninum* caused mummification.

Table 9
Seroprevalence of *N. caninum* antibodies in free range wildlife.

Host	Country	No. examined	Assay	Cut-off titer or test	% Positive	Reference
Canids						
Coyote (<i>Canis latrans</i>)	USA	12	IFAT	1:100	16.7	Stieve et al. (2010)
Gray wolf (<i>Canis lupus</i>)	Scandinavia	109	IB, ELISA	ISCOM	3.7	Björkman et al. (2010)
	Spain	28	NAT, ELISA	VMRD	21.4	Sobrino et al. (2008)
	USA Alaska	324	IFAT	1:100	9.0	Stieve et al. (2010)
	Yellow Stone Park	220	IFAT	1:50	50.0	Almberg et al. (2009)
Red fox (<i>Vulpes vulpes</i>)	Ireland	220	IFAT	1:50	3.0	Murphy et al. (2007)
	Spain	95	NAT, ELISA, IFAT	VMRD	3.2	Sobrino et al. (2008)
		53	NAT	1:40	69.8	Marco et al. (2008)
Mustelids						
Stone martin (<i>Martes foina</i>)	Spain	14	NAT, ELISA, IFAT	VMRD	21.4	Sobrino et al. (2008)
Pine martin (<i>Martes martes</i>)	Spain	3	NAT, ELISA	VMRD	66.7	Sobrino et al. (2008)
Eurasian badger (<i>Meles meles</i>)	Spain	31	NAT, ELISA	VMRD	6.4	Sobrino et al. (2008)
Pole cat (<i>Mustella putorius</i>)	Spain	2	NAT, ELISA	VMRD	50	Sobrino et al. (2008)
Felids						
Feral cat (<i>Felis silvestris</i>)	Spain	59	IFAT, ELISA	VMRD	6.8	Millán et al. (2009b)
Eurasian lynx (<i>Lynx lynx</i>)	Spain	26	ELISA	VMRD	19	Millán et al. (2009a)
Iberian lynx (<i>Lynx pardinus</i>)	Spain	25	ELISA, IFAT	VMRD	12.0	Sobrino et al. (2008)
European wild cat (<i>Felis silvestris</i>)	Spain	6	NAT, ELISA	VMRD	16.7	Sobrino et al. (2008)
Egyptian mongoose (<i>Herpestes ichneumon</i>)	Spain	21	ELISA	VMRD	13	Millán et al. (2009a)
Cervids and ruminants						
Spanish ibex (<i>Capra pyrenaica hispanica</i>)	Spain	531	ELISA, IFAT	VMRD, 1:50	5.6	García-Bocanegra et al. (in press)
Mouflon (<i>Ovis ammon</i>) ^a	Czech Republic	105	IFAT	1:50	3	Bártová et al. (2007)
Mule deer (<i>Odocoileus hemionus hemionus</i>)	USA	42	NAT	1:25	16.6	Dubey et al. (2008a)
Black-tailed deer (<i>Odocoileus hemionus columbianus</i>)	USA	43	NAT	1:25	18.6	Dubey et al. (2008a)
White-tailed deer (<i>Odocoileus virginianus</i>)	USA-Iowa	170	NAT, IFAT	1:25	88.2	Dubey et al. (2009)
	USA-Minnesota	62	NAT, IFAT	1:25	70.0	Dubey et al. (2009)
Vietnam sika deer (<i>Cervus nippon pseudaxis</i>)	Czech Republic	14	IFAT, ELISA	1:50	14.0	Bártová et al. (2007)
Roe deer (<i>Capreolus capreolus</i>)	Belgium	73	ELISA	ID-VET	2.7	de Craeye et al. (2011)
	Czech Republic ^a	79	IFAT	1:50	14.0	Bártová et al. (2007)
	Spain Galicia	160	ELISA	VMRD	13.7	Panadero et al. (2010)
	Sweden	199	ELISA, IB	SVANOVA	1	Malmsten et al. (2011)
Fallow deer (<i>Dama dama</i>)	Belgium	4	ELISA	ID-VET	0	de Craeye et al. (2011)
	Czech Republic ^a	143	IFAT	1:50	1.4	Bártová et al. (2007)
	Poland	47 feral 106 farmed 335	ELISA, IB ELISA	IH-ISCOM IH-ISCOM IDEXX	13 11 2.9	Goździk et al. (2010) Goździk et al. (2010) Bień et al. (in press)
Red deer (<i>Cervus elaphus</i>)	Belgium	7	ELISA	ID-VET	0	de Craeye et al. (2011)
Caribou (<i>Rangifer tarandus</i>)	USA-Alaska	453	IFAT	1:100	11.5	Stieve et al. (2010)
Moose (<i>Alces alces</i>)	Sweden	417	IH-ISCOM, ELISA	IH-ISCOM	1.0	Malmsten et al. (2011)
	USA-Alaska	201	IFAT	1:100	0.5	Stieve et al. (2010)
Marine mammals						
Kuril harbor seal (<i>Phoca vitulina stejnegeri</i>)	Japan	234	ELISA, IH-tNcSAG1	In house	5.9	Fujii et al. (2007)
Spotted seal (<i>Phoca largha</i>)	Japan	13	ELISA, IH-tNcSAG1	In house	15.3	Fujii et al. (2007)
North American opossum (<i>Didelphis virginiana</i>)	USA	30	IFAT	1:100	0	Houk et al. (2010)
Sea otter (<i>Enhydra lutris nerensis</i>)	USA	16	IFAT	1:40	68.7	Miller et al. (2010)
Avians						
Common raven (<i>Corvus corax</i>)	Spain	67	IFAT	1:50	35.8	Molina-López et al. (in press)

^a Some of the animals were from hunting farms.

Neosporosis-induced abortions occur year-round. In some epidemic herd outbreaks as many as 57% of dairy cows have been reported to abort over just a few weeks up to months. Abortion outbreaks have been defined as epidemic if more than 10% or 12.5% of cows at risk abort within six to eight weeks. A small proportion (<5%) of cows may

have repeated abortion due to neosporosis (Dubey et al., 2007a; Pabón et al., 2007). A previous *N. caninum* abortion is not considered a cause of sterility (Santolaria et al., 2009).

Clinical signs, other than abortion, which have only been reported in calves <two months of age, include neurologic signs, an inability to rise and below average birth

weight. The hind limbs and/or the forelimbs may be flexed or hyperextended and neurologic examination may reveal ataxia, decreased patellar reflexes, and loss of conscious proprioception. Exophthalmia or an asymmetrical appearance in the eyes may be achieved and occasionally birth defects including hydrocephalus and a narrowing of the spinal cord may occur.

3.4. Diagnosis

Diagnosis of neosporosis abortion is difficult, and often expensive. Serologic examination of the dam and the fetus, and the detection of lesions and *N. caninum* in a fetus by immunohistology and PCR can all aid diagnosis and we have reviewed these procedures in detail (Dubey and Schares, 2006). Establishing a cause-effect relationship between abortion and *N. caninum* is even more complex because asymptomatic congenital *N. caninum* infections are common and finding the presence of the parasite or parasite DNA does not mean that *N. caninum* caused the abortion. In comprehensive studies based on exclusion of all other causes of abortion and the observation of *N. caninum*-associated lesions and parasites in aborted fetuses in cattle from California and The Netherlands, about 20% of all abortions were associated with *N. caninum*. Recent reports of *N. caninum*-associated abortion are summarized in Table 6. In some of these only PCR was used for diagnosis of abortion; diagnostic rate varies with the method used (Sadrebazzaz et al., 2007; Sánchez et al., 2009).

Diagnosis of acute versus chronic *N. caninum* infection in adult cows is epidemiologically important. Avidity tests have been used to distinguish acute versus chronic phase of infection. Low avidity values are associated with acute infection (Basso et al., 2010) but the window for a diagnosis of an acute infection is short, i.e. lasts only several weeks post infection. Recently, Aguado-Martínez et al. (2008) described ELISAs based on the recombinant proteins NcGRA7 and NcSAG4 for the detection of antibodies during acute infection (tachyzoite replication), and chronic infection (bradyzoites), respectively. Detection of antibodies to both recombinant proteins may indicate reactivated infection.

Many serological tests are available for the diagnosis of bovine *N. caninum* infection (Dubey and Schares, 2006; Wapenaar et al., 2007a). It is worth noting that antibodies and white blood cell counts may fluctuate during pregnancy, even in non aborting cows (Lopez-Gatius et al., 2007a; Nogareda et al., 2007; Serrano et al., 2011), and some animals may become serologically negative (Nogareda et al., 2007; Dijkstra et al., 2008). The sub-class of antibodies detected may vary with clinical status; IgG2 antibodies dominated in almost all of the aborting cattle under examination while non-aborting cattle showed either a dominating IgG1 or a dominating IgG2 response (Almería et al., 2009a).

An ELISA based on the use of NcGRA7 protein was useful in the detection of antibodies in aborting cows versus those calving normally (Huang et al., 2007).

Further serological methods to diagnose *N. caninum* infection based on recombinant antigens are under development (Borsuk et al., 2011). Previous studies had

indicated that the detection of pregnancy-associated substances could aid diagnosis of neosporosis abortion. In one report, the detection of pregnancy-associated glycoprotein-1 (PAG-1) in plasma was not influenced by a chronic *N. caninum* infection. However, PAG-1 measurements in aborting animals could provide a useful parameter to assess the fetoplacental status independent of the *N. caninum* infection (Lopez-Gatius et al., 2007a).

3.5. Pathogenesis of abortion

Bovine neosporosis is mainly a disease of the placenta and fetus, initiated following a maternal parasitemia, triggered either as the result of a primary (exogenous) maternal infection or following recrudescence of a persistent (endogenous) infection during pregnancy (Dubey et al., 2007a). Following a parasitemia *N. caninum* is able to establish itself in the maternal caruncular septum before crossing to the fetal placental villus. For abortion to occur the fetus, or its placenta, has to be so damaged that it is no longer viable and several factors may interact, to a greater or lesser extent, to influence this (Gibney et al., 2008). Primary parasite-induced placental damage may jeopardize fetal survival directly or cause release of maternal prostaglandins that in turn cause luteolysis and abortion. Fetal damage may occur due to primary tissue damage caused by the multiplication of *N. caninum* in the fetus or due to insufficient oxygen/nutrition, secondary to placental damage. In addition, it has been suggested that maternal immune expulsion of the fetus may occur, associated with the release of maternal pro-inflammatory cytokines in the placenta or hormonal deregulation. While clearly all these proposed mechanisms are related in one way or another, one or more of them may be more important in a given instance and all may be influenced by the stage of gestation (Dubey et al., 2006; Innes, 2007; Lopez-Gatius et al., 2007b; Gibney et al., 2008; Almería et al., 2010). Extensive lesions in vital organs can directly kill the fetus (Gibney et al., 2008). The production of regulatory cytokines (such as IL-10) and inflammatory (such as gamma interferon) cytokines, and a direct injury of the fetus by tachyzoites multiplication may determine the survival or death of the fetus (Innes, 2007; Rosbottom et al., 2007, 2008, 2011; Almería et al., 2010; Almería et al., in press). High levels of prolactin in *Neospora* infected cows may have a protective effect on gestation (García-Ispierto et al., 2009). Previous studies indicated that progesterone may have a positive effect on pregnancy in cattle by modulating the Th1/Th2 driven immune responses. In this effect, an interaction of *N. caninum* and *Coxiella burnetii* has been recently described. Non-aborting cows seropositive to both, *N. caninum* and *C. burnetii* showed higher plasma progesterone levels than the remaining animals examined (García-Ispierto et al., 2010). However, an artificial progesterone supplementation of cows during pregnancy did not reduce neosporosis abortion but increased the risk of abortion in cows with high *N. caninum* antibody titres (Bech-Sabat et al., 2007).

The risk of transmission and fetal disease maybe in part related to the stage of gestation at the time of infection. The transmission rate increases with gestational age perhaps

related to placental vascularization because the placenta seems to be more permeable in the last trimester (Dubey et al., 2006, 2007a).

3.6. Risk factors

The knowledge on risk factors for herds to acquire *N. caninum* infection and *N. caninum*-associated abortion is important for the development and the implementation of measures to control bovine neosporosis. Potential risk factors have been extensively described (Dubey et al., 2007a). It is now generally accepted that the presence of farm dogs increases the chance of *N. caninum* infection in cattle (Dubey et al., 2007a; VanLeeuwen et al., 2010a). Dogs as definitive hosts increase the chance of a postnatal infection via oocyst contamination of the cattle food or environment. Specific feeding habits of farm dogs, like feeding on aborted fetuses and placenta may increase the chance of seropositivity in cattle (VanLeeuwen et al., 2010a). Differences in the farm management (e.g. feeding, pasture management, cattle density and housing) may have an influence on infection risk. In a survey of 5594 dairy and beef cows in Argentina, seropositive animals were 85% more likely to abort than seronegative and the results suggested a higher risk of seropositivity in dairy than in beef herds possibly due to not further specified differences between dairy and beef herds regarding herd management practices (Moore et al., 2009).

In chronic, congenitally infected cattle additional influences could increase the risk of neosporosis abortion. In two Holstein-Friesian dairy herds it was shown recently that in heifers and parous cows an increase in the cumulative number of days with a mean relative humidity lower than 60% during the second trimester of pregnancy was associated with a higher risk of abortion (Yániz et al., 2010). In parous cows an increased rainfall during the second trimester of gestation had the same effect (Yániz et al., 2010). The abortion risk was higher in parous cows with high *N. caninum* antibody titres and in cows which had been inseminated with Friesian semen compared to those inseminated with Limousin or Belgian Blue semen (Yániz et al., 2010).

N. caninum is considered a primary pathogen. However, concurrent infections may aggravate neosporosis. Bovine herpes virus 1 co-infection was found in 27% of 948 cattle and considered a potential risk factor for bovine neosporosis in Italy (Rinaldi et al., 2007). In a small survey in Vietnam, there was a strong association between bovine virus diarrhoea (BVD) and *Neospora* seropositivity (Duong et al., 2008). In a Canadian study, *N. caninum* seropositivity had a negative effect on reproduction parameters like first-service conception and calving intervals. Interestingly there seemed to be an interaction between BVD and neosporosis since *N. caninum* seropositivity had the negative effects on the parameter “first-service conception” mainly in BVD seronegative dams (VanLeeuwen et al., 2010b). Similar interactions were not observed regarding *Mycobacterium avium paratuberculosis* or Bovine leukemia virus seropositivity (VanLeeuwen et al., 2010b).

Another Canadian study examined the effects of climate (aridity), soil pH and agroecological region on seropositiv-

ity for *Mycobacterium avium paratuberculosis*, *N. caninum*, Bovine leukemia virus, and Bovine viral diarrhoea virus. No significant effects were observed regarding *N. caninum* (Scott et al., 2007).

3.7. Prevention and control

The major economic loss due to neosporosis is reproductive failure in cattle in many countries. In addition to the direct costs involved in fetal loss, indirect costs include professional help and expenses associated with establishing a diagnosis, rebreeding, possible loss of milk yield, and replacement costs if aborted cows are culled. The diagnosis of neosporosis-associated abortion is difficult and expensive. Postnatal losses due to neosporosis are difficult to document because there are no obvious ill effects in adult cattle, other than fetal loss. Culling perhaps accounts for the major loss associated with neosporosis. In general, less is known of the causes of abortion in beef cattle than in dairy cattle because of the difficulty of monitoring when small fetuses are expelled in the first trimester, and so there are no accurate assessments of *N. caninum*-induced losses in beef cattle.

The economic production losses, other than reproduction are difficult to assess because there are no clinical signs in adult cattle. Although neosporosis has been reported to decrease milk production and weight gain, in a recent study, no differences in weight gain was found in seropositive versus negative beef cattle (Hoar et al., 2007; Moré et al., 2010). Effects of *N. caninum* infection on milk production are not clear, yet. In a Canadian study *N. caninum* seropositivity was associated with lower production of milk, fat and protein in primiparous cows versus *N. caninum*-seronegative primiparous cows (Tiwari et al., 2007).

Many controls measures have been discussed to reduce *N. caninum* infection in cattle (Dubey et al., 2007a), including embryo transfer, artificial insemination of seropositive dams with semen from beef bulls, culling, replacement heifers, chemotherapy, and vaccination. Studies from Spain indicated that the likely hood of abortion was significantly lower for heifers and parous cows inseminated with beef bull semen compared with those inseminated with Holstein-Friesian bull semen (Almería et al., 2009b; Yániz et al., 2010); this effect might be due to a favorable effect of cross-breed pregnancies on placental function.

Transfer of embryos from infected dams into uninfected recipients can prevent endogenous transplacental transmission of *N. caninum*. Embryo transfer should only be accomplished to seronegative recipient cows. This technique may be used to recover uninfected calves from genetically valuable but *N. caninum*-infected dams. As a consequence, pre-transfer testing of recipients for infection with *N. caninum* is highly recommended (Paz et al., 2007; de Oliveira et al., 2010). Congenital infection and abortion can occur if recipient cows are seropositive (de Oliveira et al., 2010). The earliest stage of pregnancy, the fetus acquires *N. caninum* infection is unknown; embryos from seropositive cows were found to be not infected with *N. caninum* (Moskwa et al., 2008).

Annual serological screening could be useful in a control strategy because a long term-study showed that *N. caninum* seropositivity was very stable during the observation period and *N. caninum* seropositive cows showed a high rate of repeat abortions (Pabón et al., 2007). A 'test and cull' strategy may include the following options: (i) test and cull seropositive dams or seropositive aborting dams; (ii) test and inseminate the progeny of seropositive dams with beef bull semen only; and (iii) test and exclude the progeny of seropositive dams from breeding.

Mathematical models can be applied to estimate costs of bovine neosporosis and to determine the benefit and the efficiency of various control measures (Häsler et al., 2008; Reichel and Ellis, 2009). By a mathematical approach losses were estimated to be 81–1875 Euros per small farm in Switzerland (Häsler et al., 2008). A modeling of the costs and benefits of different measures to control *N. caninum* "culling of animals that had experienced *N. caninum*-associated abortion" or "not breeding replacements from *N. caninum*-associated abortion" were not cost effective. However, the measure "not breeding replacements from *N. caninum* seropositive cows" on farms with a high prevalence (50%) was financially attractive (Häsler et al., 2008).

Treatment of cattle appears to be uneconomical in cattle due to the fact that it can only be used as a preventive measure and hence it must be long-term and likely produce unacceptable milk or meat residues or withdrawal periods. Currently, there is no chemotherapy for bovine neosporosis that has been shown to be safe and effective and any effort to treat cattle with existing drugs must therefore be discouraged at this stage. However, interesting experimental studies that may result in an option for chemotherapeutic control at a later stage have been conducted. An effect of toltrazuril and its derivative ponazuril on tachyzoites on *N. caninum* has been shown *in vitro* and *in vivo* in calves (Strohbusch et al., 2009). In calves treated with ponazuril, the parasite was no longer detectable in the brain and other organs. Attempts were made to alter the course of *N. caninum* infection by prophylactic medication with a slow-release Monensin bolus in cows but results were inconclusive (VanLeeuwen et al., 2011). In an epidemiological survey, feeding monensin in dry cows in Canada had a reduced risk for *N. caninum* seropositivity (VanLeeuwen et al., 2010b).

At present there is no commercial vaccine for neosporosis. Mouse models are being used to test efficacy of killed and recombinant *N. caninum* vaccines (e.g. Aguado-Martínez et al., 2009; Rojo-Montejo et al., 2011).

Studies indicate that cattle develop cellular and humoral immunity after inoculation with killed vaccine formulations (Innes et al., 2007; Baszler et al., 2008; Moore et al., 2011). Experimentally, fetal death can be prevented in cows vaccinated with live tachyzoites in exogenously challenged cows but such protection has not been demonstrated in endogenously infected cows. As a first step towards developing a non-infectious vaccine, Baszler et al. (2008) showed that cows injected with NcSRS2 immunogens developed cellular and humoral immunity.

4. Neosporosis in dogs

Serological surveys indicate widespread exposure to the parasite worldwide (Table 3). Most canine isolates of *N. caninum* were obtained from sick dogs. Recently, *N. caninum* DNA was detected in 28 (32%) of 87 asymptomatic pound dogs in Algeria; 19 of these dogs were seronegative and 8 seropositive dogs were PCR negative (Ghalmi et al., 2008), but results needs confirmation.

N. caninum is a primary pathogen in dogs and can cause clinical disease in dogs of all ages. Most cases of clinical canine neosporosis have been in congenitally infected dogs involving littermates and previous reports were summarized earlier (Lindsay and Dubey, 2000; Dubey and Lappin, 2006). In most instances dogs are born asymptomatic and begin to develop clinical signs three or more weeks after birth. Not all pups in the litter are affected (Dubey et al., 2004, 2005, 2007c). Heckerroth and Tenter (2007) tested progeny of a Doberman bitch in Germany; three of nine pups from litter 1, one of five pups from litter 2, and one of eight pups from litter 3 were seropositive, but only one pup from litter 3 developed clinical neosporosis. The clinically affected pup had the highest IFAT titer (1:5129) and showed an unique pattern in Western blot examination.

Paralysis of rear limbs, often with contracture, is the most consistent sign of neonatal neosporosis. A wide array of clinical signs have been reported in older dogs, often on immunosuppressive therapy (Holmberg et al., 2006; Crookshanks et al., 2007; Fry et al., 2009; Galgut et al., 2010; Garosi et al., 2010). Treatment of clinical neosporosis with currently available drugs, including clindamycin, is only partially effective. None of the drugs kill *N. caninum* tissue cysts (Dubey et al., 2004, 2007c). Co-infection of *T. gondii* could occur with *N. caninum* infection, and should be considered in differential diagnosis. A bitch that had high *N. caninum* IFAT titer during pregnancy delivered pups coinfecting with *T. gondii* and *N. caninum*; the bitch was seronegative to *T. gondii* but viable *T. gondii* was isolated from the pups and *N. caninum* DNA was found in the brains of two of the six pups (Al-Qassab et al., 2009b).

5. Neosporosis in sheep and goats

The economic, clinical, and epidemiologic importance of *N. caninum* infection in sheep remains uncertain. Recent serological surveys in Table 7 indicate a very low (0.6% in New Zealand) to high (30.8% in Brazil) prevalence in asymptomatic sheep. Occasionally, neosporosis can cause abortion, neonatal mortality, and perhaps clinical signs in adult sheep. *N. caninum* DNA was detected in brains of 3 of 18 aborted fetuses from seven farms in New Zealand (Howe et al., 2008), 2% of 31 aborted fetuses in Italy (Masala et al., 2007), and in the brain of 1 of 7 sheep in Jordan (Abo-Shehada and Abu-Halaweh, 2010). West et al. (2006) found antibodies in fetuses of 4 of 5 aborted fetuses from 2 flocks in New Zealand. These studies indicate transplacental transmission of *N. caninum* in sheep but frequency and the etiology of abortion need definitive evidence. Finding of *N. caninum* DNA in the brain of an adult encephalitic Merino ewe in Australia (Bishop et al., 2010) suggests the parasite might cause clinical neosporosis in adult sheep.

Sheep are an excellent ruminant model for testing efficacies of vaccines against neosporosis abortion. Recently, Weston et al. (2009) performed an excellent dose titration of *N. caninum* tachyzoites in 90 day gestational ewes and concluded the clinical outcome was dose-dependent, ranging from 100% abortion in ewes infected with 10^6 parasites versus 50% abortion in those infected with 10^5 tachyzoites, and no abortion in those given 50 tachyzoites.

Serologic surveys indicate 2–23% prevalence in goats (Table 7). *N. caninum* DNA was found in 8.6% of 31 aborted goat fetuses in Italy (Masala et al., 2007).

6. Neosporosis in miscellaneous domestic animals

Llamas and alpacas are important to the economy of some South American countries. *N. caninum* DNA was detected in the brain or heart of 2 of 7 llama aborted fetuses, and 7 of 12 alpaca aborted fetuses from Peru (Serrano-Martínez et al., 2007a).

Although *N. caninum* has been isolated from water buffaloes from Brazil by feeding tissues of naturally-infected animals to dogs, and examining dog feces for excretion of oocysts (Neto et al., 2011), there is no report of clinical neosporosis in buffalo. However, *N. caninum* was recently found in one of nine fetuses obtained from slaughtered buffaloes in Brazil (Chryssafidis et al., 2011).

Antibodies to *N. caninum* have been found in several other domestic animals (Table 8), but clinical neosporosis has not been reported in these hosts.

7. Neosporosis in horses

Another species of *Neospora*, *N. hughesi* is considered to parasitize equids. At present it is uncertain whether *N. caninum* also infects horses because these species cross-react serologically (Gondim et al., 2009). All three viable isolates of *Neospora* from horses were identified as *N. hughesi*. Antibodies (IFAT, 1:50) to *Neospora* were reported in 11.9% of 800 horses from Israel with higher seropositivity in horses with clinical signs and in aborted mare (Kligler et al., 2007), in 24% (c-ELISA-VMRD) of 552 horses from Czech Republic (Bártová et al., 2010a), 9.3% of 57 horses from Turkey (Kilbaş et al., 2008), and in 1 of 315 horses from Costa Rica (Dangoudoubiyam et al., 2011). High level antibodies to *N. hughesi* were detected in presuckling colostrum sera of naturally exposed foals from three of 32 mares in California, USA (Pusterla et al., 2011). It is of interest that clinical neosporosis in adult horses has been reported only from the USA, including recent cases from California (Finno et al., 2007, 2010). One of these cases was in a 23 year old mule that had myeloencephalitis (Finno et al., 2010).

8. Neosporosis in wild animals

Antibodies to *N. caninum* have been found in a variety of free range wildlife (Table 9).

In addition, *N. caninum* antibodies have been found in wildlife in captivity. André et al. (2010) reported seroprevalence in following captive wildlife in zoos in Brazil: ocelot (*Leopardus pardalis*, 30 of 42), little spotted cat (*Leopardus tigrinus*, 11 of 35), Jaguar (*Panthera onca*, 8 of 13) Puma

(*Puma concolor*, 5 of 18), Jaguarandi (*Puma yagouaroundi*, 5 of 25), tiger (*Panthera tigris*, 4 of 6), Pampas cat (*Oncifelis colocolo*, 3 of 3), caracal (*Caracal caracal*, 1 of 1), serval (*Leptailurus serval*, 1 of 1), lion (*Panthera leo* 1 of 9), Fishing cat (*Prionailurus viverrinus*, 1 of 1), Bush dog (*Speothos venaticus*, 16 of 27), Crab-eating fox (*Cerdocyon thous*, 13 of 39), Maned wolf (*Cerdocyon brachyurus* 5 of 21), Hoary fox (*Pseudalopex vetulus* 4 of 7), and European wolf (*Canis lupus*, 2 of 3) tested by IFAT titer of 1:25 or higher.

Qin et al. (2007) found antibodies (IFAT, 1:40) in 3 of 73 Red pandas from zoos in People's Republic of China (PRC). Yu et al. (2009) reported 27.2% seropositivity (c-ELISA, VMRD) in 103 farm-bred blue foxes (*Alopex lagopus*) also from PRC.

Recently, *N. caninum* DNA was reported in brains of 6.6% of 304 red foxes, and 10% of 20 roe deer from Belgium (de Craeye et al., 2011). On the contrary, brain samples from 528 foxes, 224 wild mice, 16 deer and roe deer as well as from a wild boar were examined negative for the presence of *N. caninum* DNA by real time PCR (Constantin et al., 2011).

Clinical neosporosis in wildlife is rare (Dubey, 2003b). Fatal neosporosis has been diagnosed previously in a 16 day old rhinoceros, and in wild cervids (reviewed in Dubey, 2003a; Dubey et al., 2007a). Recently, fatal neosporosis was diagnosed in an aborted fetus from a white rhinoceros (*Ceratotherium simum*) in Taronga zoo Australia (Sangster et al., 2010). Hepatitis was the main lesion and the parasite was demonstrated histologically and by PCR. Both of these rhinos were originally from South Africa. A third case of acute fatal neosporosis was diagnosed in a 16-year old rhinoceros that died suddenly in captivity in a zoo in Thailand (Sommanustweechai et al., 2010). This animal also had massive hepatitis, with histologically demonstrable tachyzoites. Unlike the previous two cases, this animal was originally wild caught in Thailand.

Conflict of interest

The authors have no conflict of interest.

Acknowledgements

We would like to thank Drs. Eva Bártová, Shanti Choudhary, Leandra Ferreira, and Juliana Martins for their help.

References

- Abo-Shehadeh, M.N., Abu-Halaweh, M.M., 2010. Flock-level seroprevalence of, and risk factors for, *Neospora caninum* among sheep and goats in northern Jordan. *Prev. Vet. Med.* 93, 25–32.
- Aguado-Martínez, A., Alvarez-García, G., Fernández-García, A., Risco-Castillo, V., Arnaiz-Seco, I., Rebordosa-Trigueros, X., Navarro-Lozano, V., Ortega-Mora, L.M., 2008. Usefulness of rNcGRA7- and rNcSAG4-based ELISA tests for distinguishing primo-infection, recrudescence, and chronic bovine neosporosis. *Vet. Parasitol.* 157, 182–195.
- Aguado-Martínez, A., Alvarez-García, G., Fernández-García, A., Risco-Castillo, V., Marugán-Hernández, V., Ortega-Mora, L.M., 2009. Failure of a vaccine using immunogenic recombinant proteins rNcSAG4 and rNcGRA7 against neosporosis in mice. *Vaccine* 27, 7331–7338.
- Aguiar, D.M., Chiebao, D.P., Rodrigues, A.A.R., Cavalcante, G.T., Labruna, M.B., Gennari, S.M., 2004. Prevalência de anticorpos anti-*Neospora caninum* em ovinos do município de Monte Negro, RO Amazônia Ocidental Brasileira. *Arq. Inst. Biol. (São Paulo)* 71 supl., 616–618.

- Al-Majali, A.M., Jawasreh, K.I., Talafha, H.A., Talafha, A.Q., 2008. Neosporosis in sheep and different breeds of goats from southern Jordan: prevalence and risk factors analysis. *Am. J. Anim. Vet. Sci.* 3, 47–52.
- Al-Qassab, S., Reichel, M.P., Ivens, A., Ellis, J.T., 2009a. Genetic diversity amongst isolates of *Neospora caninum*, and the development of a multiplex assay for the detection of distinct strains. *Mol. Cell. Probes.* 23, 132–139.
- Al-Qassab, S., Reichel, M.P., Su, C., Jenkins, D., Hall, C., Windsor, P.A., Dubey, J.P., Ellis, J., 2009b. Isolation of *Toxoplasma gondii* from the brain of a dog in Australia and its biological and molecular characterization. *Vet. Parasitol.* 164, 335–339.
- Al-Qassab, S., Reichel, M.P., Ellis, J., 2010. A second generation multiplex PCR for typing strains of *Neospora caninum* using six DNA targets. *Mol. Cell. Probes.* 24, 20–26.
- Almberg, E.S., Mech, L.D., Smith, D.W., Sheldon, J.W., Crabtree, R.L., 2009. A serological survey of infectious disease in Yellowstone National Park's canid community. *PLoS ONE* 4, e7042–e7050.
- Almería, S., Nogareda, C., Santolaria, P., García-Ispuerto, I., Yáñez, J.L., López-Gatius, F., 2009a. Specific anti-*Neospora caninum* IgG1 and IgG2 antibody responses during gestation in naturally infected cattle and their relationship with gamma interferon production. *Vet. Immunol. Immunopathol.* 130, 35–42.
- Almería, S., López-Gatius, F., García-Ispuerto, I., Nogareda, C., Bech-Sabat, G., Serrano, B., Santolaria, P., Yáñez, J.L., 2009b. Effects of crossbreed pregnancies on the abortion risk of *Neospora caninum*-infected dairy cows. *Vet. Parasitol.* 163, 323–329.
- Almería, S., Araujo, R., Tuo, W., López-Gatius, F., Dubey, J.P., Gasbarre, L.C., 2010. Fetal death in cows experimentally infected with *Neospora caninum* at 110 days of gestation. *Vet. Parasitol.* 169, 304–311.
- Almería, S., Araujo, R.N., Darwich, L., Dubey, J.P., Gasbarre, L.C., 2011. Cytokine gene expression at the materno-foetal interface after experimental *Neospora caninum* infection of heifers at 110 days of gestation. *Parasite Immunol.*, in press.
- André, M.R., Adania, C.H., Teixeira, R.H.F., Silva, K.F., Jusi, M.M.G., Machado, S.T.Z., de Bortolli, C.P., Falcade, M., Sousa, L., Alegretti, S.M., Felipe, P.A.N., Machado, R.Z., 2010. Antibodies to *Toxoplasma gondii* and *Neospora caninum* in captive neotropical and exotic wild canids and felids. *J. Parasitol.* 96, 1007–1009.
- Andreotti, R., Matos, M.F.C., Gonçalves, K.N., Oshiro, L.M., Lima-Junior, M.S.C., Paiva, F., Leite, F.L., 2009. Comparison of indirect ELISA based on recombinant protein NcSRS2 and IFAT for detection of *Neospora caninum* antibodies in sheep. *Rev. Bras. Parasitol. Vet.* 18, 20–24.
- Andreotti, R., Barros, J.C., Pereira, A.R., Oshiro, L.M., Cunha, R.C., Figueiredo Neto, L.F., 2010. Association between seropositivity for *Neospora caninum* and reproductive performance of beef heifers in the Pantanal of Mato Grosso do Sul, Brazil. *Rev. Bras. Parasitol. Vet.* 19, 119–123.
- Armengol, R., Pabón, M., Santolaria, P., Cabezon, O., Adelantado, C., Yáñez, J., López-Gatius, F., Almería, S., 2007. Low seroprevalence of *Neospora caninum* infection associated with the limousin breed in cow-calf herds in Andorra, Europe. *J. Parasitol.* 93, 1029–1032.
- Azevedo, S.S., Pena, H.F., Alves, C.J., Guimaraes Filho, A.A., Oliveira, R.M., Maksimov, P., Schares, G., Gennari, S.M., 2010. Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in swine from Northeastern Brazil. *Rev. Bras. Parasitol. Vet.* 19, 80–84.
- Bandini, L.A., Neto, A.F.A., Pena, H.F.J., Cavalcante, G.T., Schares, G., Nishi, S.M., Gennari, S.M., 2011. Experimental infection of dogs (*Canis familiaris*) with sporulated oocysts of *Neospora caninum*. *Vet. Parasitol.* 176, 151–156.
- Barratt, J., Al Qassab, S., Reichel, M.P., Ellis, J.T., 2008. The development and evaluation of a nested PCR assay for detection of *Neospora caninum* and *Hammondia heydorni* in feral mouse tissues. *Mol. Cell. Probes.* 22, 228–233.
- Bartels, C.J.M., Huinink, I., Beiboer, M.L., van Schaik, G., Wouda, W., Dijkstra, T., Stegeman, A., 2007. Quantification of vertical and horizontal transmission of *Neospora caninum* infection in Dutch dairy herds. *Vet. Parasitol.* 148, 83–92.
- Basso, W., Herrmann, D.C., Conraths, F.J., Pantchev, N., Vrhovec, M.G., Schares, G., 2009a. First isolation of *Neospora caninum* from the faeces of a dog from Portugal. *Vet. Parasitol.* 159, 162–166.
- Basso, W., Schares, S., Bärwald, A., Herrmann, D.C., Conraths, F.J., Pantchev, N., Vrhovec, M.G., Schares, G., 2009b. Molecular comparison of *Neospora caninum* oocyst isolates from naturally infected dogs with cell culture-derived tachyzoites of the same isolates using nested polymerase chain reaction to amplify microsatellite markers. *Vet. Parasitol.* 160, 43–50.
- Basso, W., Schares, S., Minke, L., Bärwald, A., Maksimov, A., Peters, M., Schulze, C., Müller, M., Conraths, F.J., Schares, G., 2010. Microsatellite typing and avidity analysis suggest a common source of infection in herds with epidemic *Neospora caninum*-associated bovine abortion. *Vet. Parasitol.* 173, 24–31.
- Baszler, T.V., Shkap, V., Mwangi, W., Davies, C.J., Mathison, B.A., Mazuz, M., Resnikov, D., Fish, L., Leibovitch, B., Staska, L.M., Savitsky, I., 2008. Bovine immune response to inoculation with *Neospora caninum* surface antigen SRS2 lipopeptides mimics immune response to infection with live parasites. *Clin. Vaccine Immunol.* 15, 659–667.
- Bártová, E., Sedlák, K., Pavlík, I., Literák, I., 2007. Prevalence of *Neospora caninum* and *Toxoplasma gondii* antibodies in wild ruminants from the countryside or captivity in the Czech Republic. *J. Parasitol.* 93, 1216–1218.
- Bártová, E., Sedlák, K., Literák, I., 2009. *Toxoplasma gondii* and *Neospora caninum* antibodies in sheep in the Czech Republic. *Vet. Parasitol.* 161, 131–132.
- Bártová, E., Sedlák, K., Syrová, M., Literák, I., 2010a. *Neospora* spp. and *Toxoplasma gondii* antibodies in horses in the Czech Republic. *Parasitol. Res.* 107, 783–785.
- Bártová, E., Sedlák, K., Tremel, F., Holko, I., Literák, I., 2010b. *Neospora caninum* and *Toxoplasma gondii* antibodies in European brown hares in the Czech Republic, Slovakia and Austria. *Vet. Parasitol.* 171, 155–158.
- Bech-Sabat, G., López-Gatius, F., Santolaria, P., García-Ispuerto, I., Pabón, M., Nogareda, C., Yáñez, J.L., Almería, S., 2007. Progesterone supplementation during mid-gestation increases the risk of abortion in *Neospora*-infected dairy cows with high antibody titres. *Vet. Parasitol.* 145, 164–167.
- Benetti, A.H., Toniollo, G.H., dos Santos, T.R., Gennari, S.M., da Costa, A.J., Dias, R.A., 2008. Ocorrência de anticorpos anti-*Neospora caninum* em cães no município de Cuiabá, Mato Grosso. *Ciência Animal Bras.* 9, 177–180.
- Benetti, A.H., Schein, F.B., dos Santos, T.R., Toniollo, G.H., da Costa, A.J., Mineo, J.R., Lobato, J., de Oliveira Silva, D.A., Gennari, S.M., 2009. Pesquisa de anticorpos anti-*Neospora caninum* em bovinos leiteiros, cães e trabalhadores rurais da região Sudoeste do Estado de Mato Grosso. *Rev. Bras. Parasitol. Vet.* 18 (Suppl 1), 29–33.
- Bieñ, J., Moskwa, B., Cabaj, W., 2010. In vitro isolation and identification of the first *Neospora caninum* isolate from European bison (*Bison bonasus bonasus* L.). *Vet. Parasitol.* 173, 200–205.
- Bieñ, J., Moskwa, B., Bogdaszewski, M., Cabaj, W., 2011. Detection of specific antibodies anti-*Neospora caninum* in the fallow deer (*Dama dama*). *Res. Vet. Sci.*, in press.
- Bishop, S., King, J., Windsor, P., Reichel, M.P., Ellis, J., Slapeta, J., 2010. The first report of ovine cerebral neosporosis and evaluation of *Neospora caninum* prevalence in sheep in New South Wales. *Vet. Parasitol.* 170, 137–142.
- Björkman, C., Jakubek, E.B., Arnemo, J.M., Malmsten, J., 2010. Seroprevalence of *Neospora caninum* in gray wolves in Scandinavia. *Vet. Parasitol.* 173, 139–142.
- Boaventura, C.M., de Oliveira, V.S.F., Melo, D.P.G., Borges, L.M.F., da Silva, A.C., 2008. Prevalência de *Neospora caninum* em cães de Goiânia. *Rev. Patol. Trop.* 37, 15–22.
- Borsuk, S., Andreotti, R., Leite, F.P., da Silva Pinto, L., Simionatto, S., Hartleben, C.P., Goetze, M., Oshiro, L.M., Matos, M.D., Berne, M.E., 2011. Development of an indirect ELISA-NcSRS2 for detection of *Neospora caninum* antibodies in cattle. *Vet. Parasitol.* 177, 33–38.
- Bresciani, K.D.S., Costa, A.J., Nunes, C.M., Serrano, A.C.M., Moura, A.B., Stobbe, N.S., Perri, S.H.V., Dias, R.A., Gennari, S.M., 2007. Ocorrência de anticorpos contra *Neospora caninum* e *Toxoplasma gondii* e estudo de fatores de risco em cães de Araçatuba – SP. *Ars Veterinaria* 23, 40–46.
- Brickell, J.S., McGowan, M.M., Wathes, D.C., 2010. Association between *Neospora caninum* seropositivity and perinatal mortality in dairy heifers at first calving. *Vet. Rec.* 167, 82–85.
- Cabezón, O., Millán, J., Gomis, M., Dubey, J.P., Ferroglio, E., Almería, S., 2010. Kennel dogs as sentinels of *Leishmania infantum*, *Toxoplasma gondii*, and *Neospora caninum* in Majorca Island, Spain. *Parasitol. Res.* 107, 1505–1508.
- Campero, C.M., Pérez, A., Moore, D.P., Crudeli, G., Benitez, D., Draghi, M.G., Cano, D., Konrad, J.L., Odeón, A.C., 2007. Occurrence of antibodies against *Neospora caninum* in water buffaloes (*Bubalus bubalis*) on four ranches in Corrientes province, Argentina. *Vet. Parasitol.* 150, 155–158.
- Cardoso, J.M.S., Funada, M.R., Soares, R.M., Gennari, S.M., 2008. Perfil sorológico dos anticorpos colostrais para *Neospora caninum* em bezerros livres da infecção. *Braz. J. Vet. Res. Anim. Sci.* 45, 379–384.
- Casas, V., Chávez, G.V., Casas, A.A., Leyva, E.V., Alvarado, V.S., Serrano, A.M., Ticona, E.S., Puray Ch, D.N., 2006. Presencia de *Neospora caninum* en llamas de una empresa ganadera de la Sierra Central. *Rev. Inv. Vet. Perú.* 17, 8–13.
- Cavalcante, G.T., Monteiro, R.M., Soares, R.M., Nishi, S.M., Alves Neto, A.F., Esmerini, P.d.O., Sercundes, M.K., Martins, J., Gennari, S.M., 2011. Shed-

- ding of *Neospora caninum* oocysts by dogs fed different tissues from naturally infected cattle. *Vet. Parasitol.* 179, 220–223.
- Cedillo, C.J.R., Martínez, M.J.J., Santacruz, A.M., Banda, R.V.M., Morales, S.E., 2008. Models for experimental infection of dogs fed with tissue from fetuses and neonatal cattle naturally infected with *Neospora caninum*. *Vet. Parasitol.* 154, 151–155.
- Chanlun, A., Emanuelson, U., Frössling, J., Aiumlamai, S., Björkman, C., 2007. A longitudinal study of seroprevalence and seroconversion of *Neospora caninum* infection in dairy cattle in northeast Thailand. *Vet. Parasitol.* 146, 242–248.
- Chryssafidis, A.L., Soares, R.M., Rodrigues, A.A.R., Carvalho, N.A.T., Gennari, S.M., 2011. Evidence of congenital transmission of *Neospora caninum* in naturally infected water buffalo (*Bubalus bubalis*) fetus from Brazil. *Parasitol. Res.* 108, 741–743.
- Collantes-Fernández, E., Gómez-Bautista, M., Miró, G., Álvarez-García, G., Pereira-Bueno, J., Frisuelos, C., Ortega-Mora, L.M., 2008. Seroprevalence and risk factors associated with *Neospora caninum* infection in different dog populations in Spain. *Vet. Parasitol.* 152, 148–151.
- Constantin, E.M., Schares, G., Großmann, E., Sauter, K., Romig, T., Hartmann, S., 2011. Untersuchungen zur Rolle des Rotfuchses (*Vulpes vulpes*) als möglicher Endwirt von *Neospora caninum*. *Berl. Münch. Tierärztl. Wochenschr.* 124, 148–153.
- Costa, K.S., Santos, S.L., Uzêda, R.S., Pinheiro, A.M., Almeida, M.A.O., Araújo, F.R., McAllister, M.M., Gondim, L.F.P., 2008. Chickens (*Gallus domesticus*) are natural intermediate hosts of *Neospora caninum*. *Int. J. Parasitol.* 38, 157–159.
- Crookshanks, J.L., Taylor, S.M., Haines, D.M., Shelton, G.D., 2007. Treatment of canine pediatric *Neospora caninum* myositis following immunohistochemical identification of tachyzoites in muscle biopsies. *Can. Vet. J.* 48, 506–508.
- Cruz-Vázquez, C., Medina-Esparza, L., Marcentes, A., Morales-Salinas, E., García-Vázquez, Z., 2008. Seroepidemiological study of *Neospora caninum* infection in dogs found in dairy farms and urban areas of Aguascalientes, Mexico. *Vet. Parasitol.* 157, 139–143.
- Cunha Filho, N.A., Lucas, A.S., Pappen, F.G., Ragozo, A.M.A., Gennari, S.M., Lucia, T., Farias, N.A.R., 2008. Fatores de risco e prevalência de anticorpos anti-*Neospora caninum* em cães urbanos e rurais do Rio Grande do Sul, Brasil. *Rev. Bras. Parasitol. Vet.* 17 (Suppl 1), 301–306.
- Czopowicz, M., Kaba, J., Szalus-Jordanow, O., Nowicki, M., Witkowski, L., Frymus, T., 2011. Seroprevalence of *Toxoplasma gondii* and *Neospora caninum* infections in goats in Poland. *Vet. Parasitol.* 178, 339–341.
- Damriyasa, I.M., Schares, G., Bauer, C., 2010. Seroprevalence of antibodies to *Neospora caninum* in *Bos javanicus* ('Bali cattle') from Indonesia. *Trop. Anim. Health Prod.* 42, 95–98.
- Dangoudoubiyam, S., Oliveira, J.B., Viquez, C., Gómez-García, A., González, O., Romero, J.J., Kwok, O.C.H., Dubey, J.P., Howe, D.K., 2011. Detection of antibodies against *Sarcocystis neurona*, *Neospora* spp., and *Toxoplasma gondii* in horses from Costa Rica. *J. Parasitol.* 97, 522–524.
- Darwich, L., Cabezon, O., Echeverria, I., Pabón, M., Marco, I., Molina-López, R., Alarcia-Alejos, O., López-Gatius, F., Lavín, S., Almería, S., 2011. Presence of *Toxoplasma gondii* and *Neospora caninum* DNA in the brain of wild birds. *Vet. Parasitol.*, submitted for publication.
- de Craeye, S., Speybroeck, N., Aizenberg, D., Dardé, M.L., Collinet, F., Tavernier, P., Van, G.S., Dorny, P., Dierick, K., 2011. *Toxoplasma gondii* and *Neospora caninum* in wildlife: common parasites in Belgian foxes and Cervidae? *Vet. Parasitol.* 178, 64–69.
- de Moraes, C.C.G., Megid, J., Pituco, E.M., Okuda, L.H., Del Fava, C., de Stefano, E., Crocci, A.J., 2008. Ocorrência de anticorpos anti-*Neospora caninum* em cães da microrregião da Serra de Botucatu, Estado de São Paulo, Brasil. *Rev. Bras. Parasitol. Vet.* 17, 1–6.
- de Oliveira, V.S.F., Álvarez-García, G., Ortega-Mora, L.M., Borges, L.M.F., da Silva, A.C., 2010. Abortions in bovines and *Neospora caninum* transmission in an embryo transfer center. *Vet. Parasitol.* 173, 206–210.
- de Yaniz, M.G., Moore, D.P., Odeón, A.C., Cano, A., Cano, D.B., Leunda, M.R., Campero, C.M., 2007. Humoral immune response in pregnant heifers inoculated with *Neospora caninum* tachyzoites by conjunctival route. *Vet. Parasitol.* 148, 213–218.
- Dijkstra, T., Lam, T.J.G.M., Bartels, C.J.M., Eysker, M., Wouda, W., 2008. Natural postnatal *Neospora caninum* infection in cattle can persist and lead to endogenous transplacental infection. *Vet. Parasitol.* 152, 220–225.
- Dubey, J.P., Barr, B.C., Barta, J.R., Bjerkås, I., Björkman, C., Blagburn, B.L., Bowman, D.D., Buxton, D., Ellis, J.T., Gottstein, B., Hemphill, A., Hill, D.E., Howe, D.K., Jenkins, M.C., Kobayashi, Y., Koudela, B., Marsh, A.E., Mattsson, J.G., McAllister, M.M., Modry, D., Omata, Y., Sibley, L.D., Speer, C.A., Trees, A.J., Uggla, A., Upton, S.J., Williams, D.J.L., Lindsay, D.S., 2002. Redescription of *Neospora caninum* and its differentiation from related coccidia. *Int. J. Parasitol.* 32, 929–946.
- Dubey, J.P., 2003a. Neosporosis in cattle. *J. Parasitol.* 89 suppl., S42–S56.
- Dubey, J.P., 2003b. Review of *Neospora caninum* and neosporosis in animals. *Korean J. Parasitol.* 41, 1–16.
- Dubey, J.P., Sreekumar, C., Knickman, E., Miska, K.B., Vianna, M.C.B., Kwok, O.C.H., Hill, D.E., Jenkins, M.C., Lindsay, D.S., Greene, C.E., 2004. Biologic, morphologic, and molecular characterisation of *Neospora caninum* isolates from littermate dogs. *Int. J. Parasitol.* 34, 1157–1167.
- Dubey, J.P., Knickman, E., Greene, C.E., 2005. Neonatal *Neospora caninum* infections in dogs. *Acta Parasitol.* 50, 176–179.
- Dubey, J.P., Lappin, M.R., 2006. Toxoplasmosis and neosporosis. In: Greene, C. (Ed.), *Infectious Diseases of the Dog Cat.*, 3rd edition. Elsevier, pp. 754–775.
- Dubey, J.P., Schares, G., 2006. Diagnosis of bovine neosporosis. *Vet. Parasitol.* 140, 1–34.
- Dubey, J.P., Buxton, D., Wouda, W., 2006. Pathogenesis of bovine neosporosis. *J. Comp. Pathol.* 134, 267–289.
- Dubey, J.P., Schares, G., Ortega-Mora, L.M., 2007a. Epidemiology and control of neosporosis and *Neospora caninum*. *Clin. Microbiol. Rev.* 20, 323–367.
- Dubey, J.P., Alvarado-Esquivel, C., Liesenfeld, O., Herrera-Flores, R.G., Ramírez-Sánchez, B.E., González-Herrera, A., Martínez-García, S.A., Bandini, L.A., Kwok, O.C.H., 2007b. *Neospora caninum* and *Toxoplasma gondii* antibodies in dogs from Durango city, Mexico. *J. Parasitol.* 93, 1033–1035.
- Dubey, J.P., Vianna, M.C.B., Kwok, O.C.H., Hill, D.E., Miska, K.B., Tuo, W., Velmurgan, G.V., Conors, M., Jenkins, M.C., 2007c. Neosporosis in Beagle dogs: clinical signs, diagnosis, treatment, isolation and genetic characterization of *Neospora caninum*. *Vet. Parasitol.* 149, 158–166.
- Dubey, J.P., Mansfield, K., Hall, B., Kwok, O.C.H., Thulliez, P., 2008a. Seroprevalence of *Neospora caninum* and *Toxoplasma gondii* in black tailed deer (*Odocoileus hemionus columbianus*) and mule deer (*Odocoileus hemionus hemionus*). *Vet. Parasitol.* 156, 310–313.
- Dubey, J.P., Stone, D., Kwok, O.C.H., Sharma, R.N., 2008b. *Toxoplasma gondii* and *Neospora caninum* antibodies in dogs from Grenada, West Indies. *J. Parasitol.* 94, 750–751.
- Dubey, J.P., Jenkins, M.C., Kwok, O.C.H., Zink, R.L., Michalski, M.L., Ulrich, V., Gill, J., Carstensen, M., Thulliez, P., 2009. Seroprevalence of *Neospora caninum* and *Toxoplasma gondii* antibodies in white-tailed deer (*Odocoileus virginianus*) from Iowa and Minnesota using four serologic tests. *Vet. Parasitol.* 161, 330–334.
- Dubey, J.P., Jenkins, M.C., Rajendran, C., Miska, K., Ferreira, L.R., Martins, J., Kwok, O.C.H., Choudhary, S., 2011. Gray wolf (*Canis lupus*) is a natural definitive host for *Neospora caninum*. *Vet. Parasitol.*, in press, doi:10.1016/j.vetpar.2011.05.018.
- Duong, M.C., Alenius, S., Huong, L.T.T., Björkman, C., 2008. Prevalence of *Neospora caninum* and bovine viral diarrhoea virus in dairy cows in Southern Vietnam. *Vet. J.* 175, 390–394.
- Eiras, C., Arnaiz, I., Álvarez-García, G., Ortega-Mora, L.M., Sanjuán, M.L., Yus, E., Diéguez, F.J., 2011. *Neospora caninum* seroprevalence in dairy and beef cattle from the northwest region of Spain, Galicia. *Prev. Vet. Med.* 98, 128–132.
- Fard, S.R.N., Khalili, M., Aminzadeh, A., 2008. Prevalence of antibodies to *Neospora caninum* in cattle in Kerman province, south east Iran. *Veterinarski Arhiv.* 78, 253–259.
- Faria, E.B., Gennari, S.M., Pena, H.F.J., Athayde, A.C.R., Silva, M.L.C.R., Azevedo, S.S., 2007. Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in goats slaughtered in the public slaughterhouse of Patos city, Paraíba State Northeast region of Brazil. *Vet. Parasitol.* 149, 126–129.
- Faria, E.B., Cavalcanti, E.F.T.S.F., Medeiros, E.S., Pinheiro, J.W., Azevedo, S.S., Athayde, A.C.R., Mota, R.A., 2010. Risk factors associated with *Neospora caninum* seropositivity in sheep from the State of Alagoas, in the northeast region of Brazil. *J. Parasitol.* 96, 197–199.
- Ferre, I., Serrano-Martínez, E., Martínez, A., Osoro, K., Mateos-Sanz, A., del-Pozo, I., Aduriz, G., Tamargo, C., Hidalgo, C.O., Ortega-Mora, L.M., 2008. Effects of re-infection with *Neospora caninum* in bulls on parasite detection in semen and blood and immunological responses. *Theriogenology* 69, 905–911.
- Ferroglio, E., Pasino, M., Romano, A., Grande, D., Pregel, P., Trisciuglio, A., 2007a. Evidence of *Neospora caninum* DNA in wild rodents. *Vet. Parasitol.* 148, 346–349.
- Ferroglio, E., Pasino, M., Ronco, F., Bena, A., Trisciuglio, A., 2007b. Seroprevalence of antibodies to *Neospora caninum* in urban and rural dogs in north-west Italy. *Zoonoses Public Health.* 54, 135–139.
- Figueredo, L.A., Dantas-Torres, F., de Faria, E.B., Gondim, L.F.P., Simões-Mattos, L., Brandão-Filho, S.P., Mota, R.A., 2008. Occurrence of antibodies to *Neospora caninum* and *Toxoplasma gondii* in dogs from Pernambuco, Northeast Brazil. *Vet. Parasitol.* 157, 9–13.

- Finno, C.J., Aleman, M., Pusterla, N., 2007. Equine protozoal myeloencephalitis associated with neosporosis in 3 horses. *J. Vet. Intern. Med.* 21, 1405–1408.
- Finno, C.J., Eaton, J.S., Aleman, M., Hollingsworth, S.R., 2010. Equine protozoal myeloencephalitis due to *Neospora hughesi* and equine motor neuron disease in a mule. *Vet. Ophthalmol.* 13, 259–265.
- Fish, L., Mazuz, M., Molad, T., Savitsky, I., Shkap, V., 2007. Isolation of *Neospora caninum* from dairy zero grazing cattle in Israel. *Vet. Parasitol.* 149, 167–171.
- Fridlund-Plugge, N., Montiani-Ferreira, F., Richartz, R.R.T.B., Dal Pizzol, J., Machado, P.C., Patrício, L.F.L., Rosinelli, A.S., Locatelli-Dittrich, R., 2008. Frequency of antibodies against *Neospora caninum* in stray and domiciled dogs from urban, periurban and rural areas from Paraná State, Southern Brazil. *Rev. Bras. Parasitol. Vet.* 17, 222–226.
- Frösling, J., Nodtvedt, A., Lindberg, A., Björkman, C., 2008. Spatial analysis of *Neospora caninum* distribution in dairy cattle from Sweden. *Geospat. Health.* 3, 39–45.
- Fry, D.R., McSparran, K.D., Ellis, J.T., Harvey, C., 2009. Protozoal hepatitis associated with immunosuppressive therapy in a dog. *J. Vet. Intern. Med.* 23, 366–368.
- Fuehrer, H.P., Blöschl, I., Siehs, C., Hassl, A., 2010. Detection of *Toxoplasma gondii* *Neospora caninum*, and Encephalitozoon cuniculi in the brains of common voles (*Microtus arvalis*) and water voles (*Arvicola terrestris*) by gene amplification techniques in western Austria (Vorarlberg). *Parasitol. Res.* 107, 469–473.
- Fujii, K., Kakumoto, C., Kobayashi, M., Saito, S., Kariya, T., Watanabe, Y., Zuan, X., Igarashi, I., Suzuki, M., 2007. Seroepidemiology of *Toxoplasma gondii* and *Neospora caninum* in seals around Hokkaido, Japan. *J. Vet. Med. Sci.* 69, 393–398.
- Furuta, P.I., Mineo, T.W.P., Carrasco, A.O.T., Godoy, G.S., Pinto, A.A., Machado, R.Z., 2007. *Neospora caninum* infection in birds: experimental infections in chicken and embryonated eggs. *Parasitology* 134, 1931–1939.
- Galgut, B.I., Janardhan, K.S., Grondin, T.M., Harkin, K.R., Wight-Carter, M.T., 2010. Detection of *Neospora caninum* tachyzoites in cerebrospinal fluid of a dog following prednisone and cyclosporine therapy. *Vet. Clin. Pathol.* 39, 386–390.
- García-Ispierto, I., Lopez-Gatius, F., Almeria, S., Yaniz, J., Santolaria, P., Serrano, B., Bech-Sabat, G., Nogareda, C., Sulon, J., de Sousa, N.M., Beckers, J.F., 2009. Factors affecting plasma prolactin concentrations throughout gestation in high producing dairy cows. *Domest. Anim. Endocrinol.* 36, 57–66.
- García-Ispierto, I., Nogareda, C., Yaniz, J., Almeria, S., Martínez-Bello, D., de Sousa, N.M., Beckers, J.F., Lopez-Gatius, F., 2010. *Neospora caninum* and *Coxiella burnetii* seropositivity are related to endocrine pattern changes during gestation in lactating dairy cows. *Theriogenology* 74, 212–220.
- García-Vazquez, Z., Rosario-Cruz, R., Mejía-Estrada, F., Rodríguez-Vivas, I., Romero-Salas, D., Fernández-Ruvalcaba, M., Cruz-Vazquez, C., 2009. Seroprevalence of *Neospora caninum* antibodies in beef cattle in three southern states of Mexico. *Trop. Anim. Health Prod.* 41, 749–753.
- García-Bocanegra, I., Cabezón, O., Pabón, M., Gómez-Guillamón, F., Arenas, A., Alcaide, E., Salas-Vega, R., Dubey, J.P., Almería, S., 2011. Prevalence of *Toxoplasma gondii* and *Neospora caninum* antibodies in Spanish ibex (*Capra pyrenaica hispanica*). *Vet. J.*, in press, doi:10.1016/j.tvjl.2010.11.011.
- García-Melo, D.P., Regidor-Cerrillo, J., Ortega-Mora, L.M., Collantes-Fernández, E., de Oliveira, V.S.F., de Oliveira, M.A.P., da Silva, A.C., 2009. Isolation and biological characterisation of a new isolate of *Neospora caninum* from an asymptomatic calf in Brazil. *Acta Parasitol.* 54, 180–185.
- García-Melo, D.P., Regidor-Cerrillo, J., Collantes-Fernández, E., Aguado-Martínez, A., Del Pozo, I., Minguijón, E., Gómez-Bautista, M., Aduriz, G., Ortega-Mora, L.M., 2010. Pathogenic characterization in mice of *Neospora caninum* isolates obtained from asymptomatic calves. *Parasitology* 137, 1057–1068.
- Garosi, L., Dawson, A., Couturier, J., Matiassek, L., de Stefani, A., Davies, E., Jeffery, N., Smith, P., 2010. Necrotizing cerebellitis and cerebellar atrophy caused by *Neospora caninum* infection: magnetic resonance imaging and clinicopathologic findings in seven dogs. *J. Vet. Intern. Med.* 24, 571–578.
- Gavrea, R.R., Cozma, V., 2010. Seroprevalence of *Neospora caninum* in cows with reproductive failure in center and northwest of Romania. *Sci. Parasitol.* 11, 67–70.
- Geurden, T., Somers, R., Thanh, N.T.G., Vien, L.V., Nga, V.T., Giang, H.H., Dorny, P., Giau, H.K., Vercruyse, J., 2008. Parasitic infections in dairy cattle around Hanoi, northern Vietnam. *Vet. Parasitol.* 153, 384–388.
- Ghalimi, F., China, B., Kaidi, R., Daube, G., Losson, B., 2008. Detection of *Neospora caninum* in dog organs using real time PCR systems. *Vet. Parasitol.* 155, 161–167.
- Ghalimi, F., China, B., Kaidi, R., Losson, B., 2009a. First epidemiological study on exposure to *Neospora caninum* in different canine populations in the Algiers District (Algeria). *Parasitol. Int.* 58, 444–450.
- Ghalimi, F., China, B., Kaidi, R., Losson, B., 2009b. Evaluation of a SRS2 sandwich commercial enzyme-linked immunosorbent assay for the detection of anti-*Neospora caninum* antibodies in bovine and canine sera. *J. Vet. Diagn. Invest.* 21, 108–111.
- Ghanem, M.E., Suzuki, T., Akita, M., Nishibori, M., 2009. *Neospora caninum* and complex vertebral malformation as possible causes of bovine fetal mummification. *Can. Vet. J.* 50, 389–392.
- Gibney, E.H., Kipar, A., Rosbottom, A., Guy, C.S., Smith, R.F., Hetzel, U., Trees, A.J., Williams, D.J.L., 2008. The extent of parasite-associated necrosis in the placenta and foetal tissues of cattle following *Neospora caninum* infection in early and late gestation correlates with foetal death. *Int. J. Parasitol.* 38, 579–588.
- Gondim, L.F.P., McAllister, M.M., Pitt, W.C., Zemlicka, D.E., 2004. Coyotes (*Canis latrans*) are definitive hosts of *Neospora caninum*. *Int. J. Parasitol.* 34, 159–161.
- Gondim, L.F.P., Lindsay, D.S., McAllister, M.M., 2009. Canine and bovine *Neospora caninum* control sera examined for cross-reactivity using *Neospora caninum* and *Neospora hughesi* indirect fluorescent antibody tests. *J. Parasitol.* 95, 86–88.
- Gondim, L.S.Q., Abe-Sandes, K., Uzêda, R.S., Silva, M.S.A., Santos, S.L., Mota, R.A., Vilela, S.M.O., Gondim, L.F.P., 2010. *Toxoplasma gondii* and *Neospora caninum* in sparrows (*Passer domesticus*) in the Northeast of Brazil. *Vet. Parasitol.* 168, 121–124.
- González-Warleta, M., Castro-Hermida, J.A., Carro-Corral, C., Cortizo-Mella, J., Mezo, M., 2008. Epidemiology of neosporosis in dairy cattle in Galicia (NW Spain). *Parasitol. Res.* 102, 243–249.
- Goździk, K., Cabaj, W., 2007. Characterization of the first Polish isolate of *Neospora caninum* from cattle. *Acta Parasitol.* 52, 295–297.
- Goździk, K., Jakubek, E.B., Björkman, C., Bien, J., Moskwa, B., Cabaj, W., 2010. Seroprevalence of *Neospora caninum* in free living and farmed red deer (*Cervus elaphus*) in Poland. *Pol. J. Vet. Sci.* 13, 117–120.
- Goździk, K., Wrzesien, R., Wielgosz-Ostolska, A., Bien, J., Kozak-Ljunggren, M., Cabaj, W., 2011. Prevalence of antibodies against *Neospora caninum* in dogs from urban areas in Central Poland. *Parasitol. Res.* 108, 991–996.
- Guedes, M.H.P., Guimarães, A.M., Rocha, C.M.B.M., Hirsch, C., 2008. Frequência de anticorpos anti-*Neospora caninum* em vacas e fetos provenientes de municípios do sul de Minas Gerais. *Rev. Bras. Parasitol. Vet.* 17, 189–194.
- Guimarães, A.M., Rocha, C.M., Oliveira, T.M., Rosado, I.R., Morais, L.G., Santos, R.R., 2009. Fatores associados à soropositividade para *Babesia*, *Toxoplasma*, *Neospora* e *Leishmania* em cães atendidos em nove clínicas veterinárias do município de Lavras MG. *Rev. Bras. Parasitol. Vet.* 18 (Suppl 1), 49–53.
- Haddadzadeh, H.R., Sadrebazzaz, A., Malmasi, A., Ardakani, H.T., Nia, P.K., Sadreshirazi, N., 2007. Seroprevalence of *Neospora caninum* infection in dogs from rural and urban environments in Tehran, Iran. *Parasitol. Res.* 101, 1563–1565.
- Hajikolaei, M.R.H., Goraninejad, S., Hamidinejat, H., Ghorbanpour, M., Paryab, R., 2007. Occurrence of *Neospora caninum* antibodies in water buffaloes (*Bubalus bulalis*) from the south-western region of Iran. *Bull. Vet. Inst. Pulawy* 51, 233–235.
- Häsler, B., Stärk, K., Gottstein, B., Reist, M., 2008. Epidemiologische und finanzielle Entscheidungsgrundlagen zur Kontrolle von *Neospora caninum* auf Schweizer Milchviehbetrieben. *Schweiz. Arch. Tierheilkd.* 150, 273–280.
- Heckerth, A.R., Tenter, A.M., 2007. Immunoanalysis of three litters born to a Doberman bitch infected with *Neospora caninum*. *Parasitol. Res.* 100, 837–846.
- Hoar, B.R., McQuarry, A.C., Hietala, S.K., 2007. Prevalence of *Neospora caninum* and persistent infection with bovine viral diarrhoea virus in dairy-breed steers in a feedlot. *J. Am. Vet. Med. Assoc.* 230, 1038–1043.
- Holmberg, T.A., Vernau, W., Melli, A.C., Conrad, P.A., 2006. *Neospora caninum* associated with septic peritonitis in an adult dog. *Vet. Clin. Pathol.* 35, 235–238.
- Hornok, S., Edelhofer, R., Joachim, A., Farkas, R., Berta, K., Repasi, A., Lakatos, B., 2008. Seroprevalence of *Toxoplasma gondii* and *Neospora caninum* infection of cats in Hungary. *Acta Vet. Hung.* 56, 81–88.
- Hosseinejad, M., Hosseini, F., Mosharrar, M., Shahbaz, S., Mahzounieh, M., Schares, G., 2010. Development of an indirect ELISA test using an affinity purified surface antigen (P38) for sero-diagnosis of canine *Neospora caninum* infection. *Vet. Parasitol.* 171, 337–342.

- Houk, A.E., Goodwin, D.G., Zajac, A.M., Barr, S.C., Dubey, J.P., Lindsay, D.S., 2010. Prevalence of antibodies to *Trypanosoma cruzi*, *Toxoplasma gondii*, *Encephalitozoon cuniculi*, *Sarcocystis neurona*, *Besnoitia darlingi*, and *Neospora caninum* in North American opossums, *Didelphis virginiana*, from Southern Louisiana. *J. Parasitol.* 96, 1119–1122.
- Howe, L., West, D.M., Collett, M.G., Tattersfield, G., Pattison, R.S., Pomroy, W.E., Kenyon, P.R., Morris, S.T., Williamson, N.B., 2008. The role of *Neospora caninum* in three cases of unexplained ewe abortions in the southern North Island of New Zealand. *Small Rumin. Res.* 75, 115–122.
- Huang, P., Liao, M., Zhang, H., Lee, E.G., Nishikawa, Y., Xuan, X., 2007. Dense-granule protein NcGRA7, a new marker for the serodiagnosis of *Neospora caninum* infection in aborting cows. *Clin. Vaccine Immunol.* 14, 1640–1643.
- Hughes, J.M., Thomasson, D., Craig, P.S., Georgin, S., Pickles, A., Hide, G., 2008. *Neospora caninum*: detection in wild rabbits and investigation of co-infection with *Toxoplasma gondii* by PCR analysis. *Exp. Parasitol.* 120, 255–260.
- Ibrahim, H.M., Huang, P., Salem, T.A., Talaat, R.M., Nasr, M.I., Xuan, X., Nishikawa, Y., 2009. Short report: prevalence of *Neospora caninum* and *Toxoplasma gondii* antibodies in northern Egypt. *Am. J. Trop. Med. Hyg.* 80, 263–267.
- Innes, E.A., 2007. The host-parasite relationship in pregnant cattle infected with *Neospora caninum*. *Parasitology* 134, 1903–1910.
- Innes, E.A., Bartley, P.M., Maley, S.W., Wright, S.E., Buxton, D., 2007. Comparative host-parasite relationships in ovine toxoplasmosis and bovine neosporosis and strategies for vaccination. *Vaccine* 25, 5495–5503.
- Jesus, E.E.V., Almeida, M.A.O., Atta, A.M., 2007. Anti-neosporal IgG and IgE antibodies in canine neosporosis. *Zoonoses Public Health* 54, 387–392.
- Kamga-Waladjo, A.R., Chatagnon, G., Bakou, S.N., Boly, H., Diop, P.E.H., Tainturier, D., 2009. *Neospora caninum* antibodies and its consequences for reproductive characteristics in wandering sows from Senegal West Africa. *Asian J. Anim. Vet. Adv.* 4, 263–266.
- Kamga-Waladjo, A.R., Gbati, O.B., Kone, P., Lapo, R.A., Chatagnon, G., Bakou, S.N., Pangu, L.J., Diop, P.H., Akakpo, J.A., Tainturier, D., 2010. Seroprevalence of *Neospora caninum* antibodies and its consequences for reproductive parameters in dairy cows from Dakar-Senegal West Africa. *Trop. Anim Health Prod.* 42, 953–959.
- Kilbaş, Z.G., Adanir, R., Avcioglu, H., 2008. Seroprevalence of *Neospora caninum* in racehorses in Ankara Turkey. *Acta Parasitol.* 53, 315–316.
- King, J.S., Slapeta, J., Jenkins, D.J., Al-Qassab, S.E., Ellis, J.T., Windsor, P.A., 2010. Australian dingoes are definitive host of *Neospora caninum*. *Int. J. Parasitol.* 40, 945–950.
- Klevar, S., Norström, M., Tharaldsen, J., Clausen, T., Björkman, C., 2010. The prevalence and spatial clustering of *Neospora caninum* in dairy herds in Norway. *Vet. Parasitol.* 170, 153–157.
- Kligler, E.B., Shkap, V., Baneth, G., Mildenberg, Z., Steinman, A., 2007. Seroprevalence of *Neospora* spp. among asymptomatic horses, aborted mares and horses demonstrating neurological signs in Israel. *Vet. Parasitol.* 148, 109–113.
- Konnai, S., Mingala, C.N., Sato, M., Abes, N.S., Venturina, F.A., Gutierrez, C.A., Sano, T., Omata, Y., Cruz, L.C., Onuma, M., Ohashi, K., 2008. A survey of abortifacient infectious agents in livestock in Luzon, the Philippines, with emphasis on the situation in a cattle herd with abortion problems. *Acta Tropica.* 105, 269–273.
- Kubota, N., Sakata, Y., Miyazaki, N., Itamoto, K., Bannai, H., Nishikawa, Y., Xuan, X., Inokuma, H., 2008. Serological survey of *Neospora caninum* infection among dogs in Japan through species-specific ELISA. *J. Vet. Med. Sci.* 70, 869–872.
- Kul, O., Kabakci, N., Yildiz, K., Ocal, N., Kalender, H., Ilkme, N.A., 2009. *Neospora caninum* associated with epidemic abortions in dairy cattle: the first clinical neosporosis report in Turkey. *Vet. Parasitol.* 159, 69–72.
- Langoni, H., Greca, H.J., Guimaraes, F.F., Ullmann, L.S., Gaio, F.C., Uehara, R.S., Rosa, E.P., Amorim, R.M., da Silva, R.C., 2011. Serological profile of *Toxoplasma gondii* and *Neospora caninum* infection in commercial sheep from Sao Paulo State, Brazil. *Vet. Parasitol.* 177, 50–54.
- Lassen, B., Viltrop, A., Raaperi, K., Jarvis, T., 2008. *Neospora caninum* antibodies in bulk milk and serum from Estonian dairy farms. In: Alitalo, I., Jarvis, T., Zilinskas, H., Bizokas, V., Steinlechner, S., Birgele, E. (Eds.), *Proc. Int. Sci. Conf. Animal Health. Food Hygiene, Jelgava, Latvia*. November 14, 2008, 105–107.
- Lima, J.T.R., Ahid, S.M.M., Barrêto, R.A., Pena, H.F.J., Dias, R.A., Gennari, S.M., 2008. Prevalência de anticorpos anti-*Toxoplasma gondii* e anti-*Neospora caninum* em rebanhos caprinos do município de Mossoró, Rio Grande do Norte. *Braz. J. Vet. Res. Anim. Sci.* 45, 81–86.
- Lindsay, D.S., Dubey, J.P., 2000. Canine neosporosis. *J. Vet. Parasitol.* 14, 1–11.
- Liu, J., Yu, J., Wang, M., Liu, Q., Zhang, W., Deng, C., Ding, J., 2007. Serodiagnosis of *Neospora caninum* infection in cattle using a recombinant tNcSRS2 protein-based ELISA. *Vet. Parasitol.* 143, 358–363.
- Liu, J., Cai, J.Z., Zhang, W., Liu, Q., Chen, D., Han, J.P., Liu, Q.R., 2008. Seroprevalence of *Neospora caninum* and *Toxoplasma gondii* infection in yaks (*Bos grunniens*) in Qinghai, China. *Vet. Parasitol.* 152, 330–332.
- Loobuyck, M., Frossling, J., Lindberg, A., Björkman, C., 2009. Seroprevalence and spatial distribution of *Neospora caninum* in a population of beef cattle. *Prev. Vet. Med.* 92, 116–122.
- Lopes, M.G., Mendonça, I.L., Fortes, K.P., Amaku, M., Pena, H.F.J., Gennari, S.M., 2011. Presence of antibodies against *Toxoplasma gondii*, *Neospora caninum* and *Leishmania infantum* in dogs from Piauí. *Rev. Bras. Parasitol.* 20, 1–4.
- Lopez-Gatius, F., Garbayo, J.M., Santolaria, P., Yaniz, J.L., Almeria, S., Ayad, A., de Sousa, N.M., Beckers, J.F., 2007a. Plasma pregnancy-associated glycoprotein-1 (PAG-1) concentrations during gestation in *Neospora*-infected dairy cows. *Theriogenology* 67, 502–508.
- Lopez-Gatius, F., Almeria, S., Donofrio, G., Nogareda, C., Garcia-Ispuerto, I., Bech-Sabat, G., Santolaria, P., Yaniz, J.L., Pabon, M., de Sousa, N.M., Beckers, J.F., 2007b. Protection against abortion linked to gamma interferon production in pregnant dairy cows naturally infected with *Neospora caninum*. *Theriogenology* 68, 1067–1073.
- Lu, Y., Wang, G., Ma, L., 2007. Serodiagnosis of *Neospora caninum* infection in cashmere goat by enzyme-linked immunosorbent assay with recombinant truncated NcSAG1t. *Chin. Anim. Husb. Vet. Med.* 34, 109–111.
- Malmsten, J., Jakubek, E.B., Björkman, C., 2011. Prevalence of antibodies against *Toxoplasma gondii* and *Neospora caninum* in moose (*Alces alces*) and roe deer (*Capreolus capreolus*) in Sweden. *Vet. Parasitol.* 177, 275–280.
- Mansourian, M., Khodakaram-Tafti, A., Namavari, M., 2009. Histopathological and clinical investigations in *Neospora caninum* experimentally infected broiler chicken embryonated eggs. *Vet. Parasitol.* 166, 185–190.
- Marco, I., Ferroglio, E., López-Olvera, J.R., Montané, J., Lavín, S., 2008. High seroprevalence of *Neospora caninum* in the red fox (*Vulpes vulpes*) in the Pyrenees (NE Spain). *Vet. Parasitol.* 152, 321–324.
- Marques, F.A.C., Headley, A.S., Figueredo-Pereira, V., Taroda, A., Barros, L.D., Cunha, I.A.L., Munhoz, K., Bugni, F.M., Zulpo, D.L., Igarashi, M., Vidotto, O., Junior, J.S., Garcia, J.L., 2011. *Neospora caninum*: evaluation of vertical transmission in slaughtered beef cows (*Bos indicus*). *Parasitol. Res.* 108, 1015–1019.
- Martins, J., Kwok, O.C.H., Dubey, J.P., 2011. Seroprevalence of *Neospora caninum* in free-range chickens (*Gallus domesticus*) from the Americas. *Vet. Parasitol.*, in press, doi:10.1016/j.vetpar.2011.05.023.
- Masala, G., Porcu, R., Daga, C., Denti, S., Canu, G., Patta, C., Tola, S., 2007. Detection of pathogens in ovine and caprine abortion samples from Sardinia Italy, by PCR. *J. Vet. Diagn. Invest.* 19, 96–98.
- McAllister, M.M., Dubey, J.P., Lindsay, D.S., Jolley, W.R., Wills, R.A., McGuire, A.M., 1998. Dogs are definitive hosts of *Neospora caninum*. *Int. J. Parasitol.* 28, 1473–1478.
- McCann, C.M., McAllister, M.M., Gondim, L.F.P., Smith, R.F., Cripps, P.J., Kipar, A., Williams, D.J.L., Trees, A.J., 2007. *Neospora caninum* in cattle: experimental infection with oocysts can result in exogenous transplacental infection, but not endogenous transplacental infection in the subsequent pregnancy. *Int. J. Parasitol.* 37, 1631–1639.
- McCann, C.M., Vyse, A.J., Salmon, R.L., Thomas, D., Williams, D.J.L., McGarry, J.W., Pebody, R., Trees, A.J., 2008. Lack of serologic evidence of *Neospora caninum* in humans, England. *Emerg. Infect. Dis.* 14, 978–980.
- Millán, J., Candela, M.G., Palomares, F., Cubero, M.J., Rodriguez, A., Baral, M., de la Fuente, J., Almeria, S., Leon-Vizcaino, L., 2009a. Disease threats to the endangered Iberian lynx (*Lynx pardinus*). *Vet. J.* 182, 114–124.
- Millán, J., Cabezón, O., Pabón, M., Dubey, J.P., Almeria, S., 2009b. Seroprevalence of *Toxoplasma gondii* and *Neospora caninum* in feral cats (*Felis silvestris catus*) in Majorca, Balearic Islands, Spain. *Vet. Parasitol.* 165, 323–326.
- Miller, M.A., Conrad, P.A., Harris, M., Hatfield, B., Langlois, G., Jessup, D.A., Magargal, S.L., Packham, A.E., Toy-Choutka, S., Melli, A.C., Murray, M.A., Gulland, F.M., Grigg, M.E., 2010. A protozoal-associated epizootic impacting marine wildlife: Mass-mortality of southern seal otters (*Enhydra lutris nereis*) due to *Sarcocystis neurone* infection. *Vet. Parasitol.* 172, 183–194.
- Mineo, T.W.P., Carrasco, A.O.T., Marciano, J.A., Werther, K., Pinto, A.A., Machado, R.Z., 2009. Pigeons (*Columba livia*) are a suitable experimental model for *Neospora caninum* infection in birds. *Vet. Parasitol.* 159, 149–153.

- Minervino, A.H.H., Ragozo, A.M.A., Monteiro, R.M., Ortolani, E.L., Gennari, S.M., 2008. Prevalence of *Neospora caninum* antibodies in cattle from Santarém, Pará, Brazil. *Res. Vet. Sci.* 84, 254–256.
- Modolo, J.R., Stachissini, A.V.M., Gennari, S.M., Dubey, J.P., Langoni, H., Padovani, C.R., Barrozo, L.V., Leite, B.L.S., 2008. Freqüência de anticorpos anti-*Neospora caninum* em soros de caprinos do estado de São Paulo e sua relação com o manejo dos animais. *Pesq. Vet. Bras.* 28, 597–600.
- Molina-López, R., Cabezon, O., Pabón, M., Darwich, L., Obón, E., Lopez-Gatius, F., Dubey, J.P., Almería, S., 2011. High seroprevalence of *Toxoplasma gondii* and *Neospora caninum* in the common raven (*Corvus corax*) in the Northeast of Spain. *Res. Vet. Sci.*, in press, doi:10.1016/j.rvsc.2011.05.011.
- Moore, D.P., deYaniz, M.G., Odeón, A.C., Cano, D., Leunda, M.R., Späth, E.A.J., Campero, C.M., 2007. Serological evidence of *Neospora caninum* infections in goats from La Rioja province, Argentina. *Small Rum. Res.* 73, 256–258.
- Moore, D.P., Regidor-Cerrillo, J., Morrell, E., Poso, M.A., Cano, D.B., Leunda, M.R., Linschinky, L., Odeón, A.C., Odriozola, E., Ortega-Mora, L.M., Campero, C.M., 2008. The role of *Neospora caninum* and *Toxoplasma gondii* in spontaneous bovine abortion in Argentina. *Vet. Parasitol.* 156, 163–167.
- Moore, D.P., Pérez, A., Agliano, S., Brace, M., Cantón, G., Cano, D., Leunda, M.R., Odeón, A.C., Odriozola, E., Campero, C.M., 2009. Risk factors associated with *Neospora caninum* infections in cattle in Argentina. *Vet. Parasitol.* 161, 122–125.
- Moore, D.P., Echaide, I., Verna, A.E., Leunda, M.R., Cano, A., Pereyra, S., Zamorano, P.I., Odeón, A.C., Campero, C.M., 2011. Immune response to *Neospora caninum* native antigens formulated with immune stimulating complexes in calves. *Vet. Parasitol.* 175, 245–251.
- Moré, G., Basso, W., Bacigalupe, D., Venturini, M.C., Venturini, L., 2008a. Diagnosis of *Sarcocystis cruzi*, *Neospora caninum*, and *Toxoplasma gondii* infections in cattle. *Parasitol. Res.* 102, 671–675.
- Moré, G., Pardini, L., Basso, W., Marín, R., Bacigalupe, D., Auad, G., Venturini, L., Venturini, M.C., 2008b. Seroprevalence of *Neospora caninum*, *Toxoplasma gondii* and *Sarcocystis* sp. in llamas (*Lama glama*) from Jujuy, Argentina. *Vet. Parasitol.* 155, 158–160.
- Moré, G., Bacigalupe, D., Basso, W., Rambeaud, M., Beltrame, F., Ramirez, B., Venturini, M.C., Venturini, L., 2009. Frequency of horizontal and vertical transmission for *Sarcocystis cruzi* and *Neospora caninum* in dairy cattle. *Vet. Parasitol.* 160, 51–54.
- Moré, G., Bacigalupe, D., Basso, W., Rambeaud, M., Venturini, M.C., Venturini, L., 2010. Serologic profiles for *Sarcocystis* sp. and *Neospora caninum* and productive performance in naturally infected beef calves. *Parasitol. Res.* 106, 689–693.
- Moskwa, B., Goździk, K., Bieñ, J., Cabaj, W., 2008. Studies on *Neospora caninum* DNA detection in the oocytes and embryos collected from infected cows. *Vet. Parasitol.* 158, 370–375.
- Munhoz, A.D., Pereira, M.J.S., Flausino, W., Lopes, C.W.G., 2009. *Neospora caninum* seropositivity in cattle breeds in the South Fluminense Paraíba Valley, state of Rio de Janeiro. *Pesq. Vet. Bras.* 29, 29–32.
- Murphy, T.M., Walochnik, J., Hassl, A., Moriarty, J., Mooney, J., Toolan, D., Sanchez-Miguel, C., O'Loughlin, A., McAuliffe, A., 2007. Study on the prevalence of *Toxoplasma gondii* and *Neospora caninum* and molecular evidence of *Encephalitozoon cuniculi* and *Encephalitozoon (Septata) intestinalis* infections in red foxes (*Vulpes vulpes*) in rural Ireland. *Vet. Parasitol.* 146, 227–234.
- Nasir, A., Ashraf, M., Khan, M.S., Yaqub, T., Javeed, A., Avais, M., Akhtar, F., 2011. Seroprevalence of *Neospora caninum* in dairy buffalo in Lahore District. *Pakistan. J. Parasitol.*, 541–543.
- Neto, A.F.A., Bandini, L.A., Nishi, S.M., Soares, R.M., Driemeier, D., Antoniassi, N.A.B., Schares, G., Gennari, S.M., 2011. Viability of sporulated oocysts of *Neospora caninum* after exposure to different physical and chemical treatments. *J. Parasitol.* 97, 135–139.
- Nogareda, C., López-Gatius, F., Santolaria, P., García-Ispuerto, I., Bech-Sàbat, G., Pabón, M., Mezo, M., Gonzalez-Warleta, M., Castro-Hermida, J.A., Yáñez, J., Almería, S., 2007. Dynamics of anti-*Neospora caninum* antibodies during gestation in chronically infected dairy cows. *Vet. Parasitol.* 148, 193–199.
- Oshiro, L.M., Matos, M.F.C., de Oliveira, J.M., Monteiro, L.A.R.C., Andreotti, R., 2007. Prevalence of anti-*Neospora caninum* antibodies in cattle from the state of Mato Grosso do Sul, Brazil. *Rev. Bras. Parasitol. Vet.* 16, 133–138.
- Osoero, K., Ortega-Mora, L.M., Martínez, A., Serrano-Martínez, E., Ferre, I., 2009. Natural breeding with bulls experimentally infected with *Neospora caninum* failed to induce seroconversion in dams. *Theriogenology* 71, 639–642.
- Pabón, M., López-Gatius, F., García-Ispuerto, I., Bech-Sàbat, G., Nogareda, C., Almería, S., 2007. Chronic *Neospora caninum* infection and repeat abortion in dairy cows: a 3-year study. *Vet. Parasitol.* 147, 40–46.
- Palavicini, P., Romero, J.J., Dolz, G., Jiménez, A.E., Hill, D.E., Dubey, J.P., 2007. Fecal and serological survey of *Neospora caninum* in farm dogs in Costa Rica. *Vet. Parasitol.* 149, 265–270.
- Panadero, R., Panceira, A., López, C., Vázquez, L., Paz, A., Díaz, P., Dacal, V., Cienfuegos, S., Fernández, G., Lago, N., Díez-Baños, P., Morrondo, P., 2010. Seroprevalence of *Toxoplasma gondii* and *Neospora caninum* in wild and domestic ruminants sharing pastures in Galicia (Northwest Spain). *Res. Vet. Sci.* 88, 111–115.
- Paradies, P., Capelli, G., Testini, G., Cantacessi, C., Trees, A.J., Otranto, D., 2007. Risk factors for canine neosporosis in farm and kennel dogs in southern Italy. *Vet. Parasitol.* 145, 240–244.
- Paz, G.F., Leite, R.C., Rocha, M.A., 2007. Associação entre sorologia para *Neospora caninum* e taxa de prehez em vacas receptoras de embriões. *Arq. Bras. Med. Vet. Zootec.* 59, 1323–1325.
- Pedraza-Díaz, S., Marugán-Hernández, V., Collantes-Fernández, E., Regidor-Cerrillo, J., Rojo-Montejo, S., Gómez-Bautista, M., Ortega-Mora, L.M., 2009. Microsatellite markers for the molecular characterization of *Neospora caninum*: application to clinical samples. *Vet. Parasitol.* 166, 38–46.
- Pescador, C.A., Corbellini, L.G., Oliveira, E.C., Raymundo, D.L., Driemeier, D., 2007. Histopathological and immunohistochemical aspects of *Neospora caninum* diagnosis in bovine aborted fetuses. *Vet. Parasitol.* 150, 159–163.
- Ploneczka, K., Mazurkiewicz, M., 2008. Seroprevalence of *Neospora caninum* in dogs in south-western Poland. *Vet. Parasitol.* 153, 168–171.
- Puray, N.C., Chávez, A.V., Casas, E.A., Falcón, P.N., Casas, G.V., 2006. Prevalencia de *Neospora caninum* en bovinos de una empresa ganaderade la sierra central del Perú. *Rev. Inv. Vet. Perú* 17, 189–194.
- Pusterla, N., Conrad, P.A., Packham, A.E., Mapes, S.M., Finno, C.J., Gardner, I.A., Barr, B.C., Ferraro, G.L., Wilson, W.D., 2011. Endogenous transplacental transmission of *Neospora hughesi* in naturally infected horses. *J. Parasitol.* 97, 281–285.
- Qin, Q., Wei, F., Li, M., Dubovi, E.J., Loeffler, I.K., 2007. Serosurvey of infectious disease agents of carnivores in captive red pandas (*Ailurus fulgens*) in China. *J. Zoo Wildlife Med.* 38, 42–50.
- Rajkhowa, S., Rajkhowa, C., Dutta, P.R., Michui, P., Das, R., 2008. Serological evidence of *Neospora caninum* infection in mithun (*Bos frontalis*) from India. *Res. Vet. Sci.* 84, 250–253.
- Razmi, G., 2009. Fecal and molecular survey of *Neospora caninum* in farm and household dogs in Mashhad Area, Khorasan Province, Iran. *Korean J. Parasitol.* 47, 417–420.
- Razmi, G.R., Maleki, M., Farzaneh, N., Talebkhani Garoussi, M., Fallah, A.H., 2007. First report of *Neospora caninum*-associated bovine abortion in Mashhad area, Iran. *Parasitol. Res.* 100, 755–757.
- Razmi, G.R., Zarea, H., Naseri, Z., 2010. A survey of *Neospora caninum*-associated bovine abortion in large dairy farms of Mashhad, Iran. *Parasitol. Res.* 106, 1419–1423.
- Regidor-Cerrillo, J., Pedraza-Díaz, S., Gómez-Bautista, M., Ortega-Mora, L.M., 2006. Multilocus microsatellite analysis reveals extensive genetic diversity in *Neospora caninum*. *J. Parasitol.* 92, 517–524.
- Regidor-Cerrillo, J., Gómez-Bautista, M., Pereira-Bueno, J., Aduriz, G., Navarro-Lozano, V., Risco-Castillo, V., Fernández-García, A., Pedraza-Díaz, S., Ortega-Mora, L.M., 2008. Isolation and genetic characterization of *Neospora caninum* from asymptomatic calves in Spain. *Parasitology* 135, 1651–1659.
- Regidor-Cerrillo, J., Pedraza-Díaz, S., Rojo-Montejo, S., Vazquez-Moreno, E., Arnaiz, I., Gomez-Bautista, M., Jimenez-Palacios, S., Ortega-Mora, L.M., Collantes-Fernandez, E., 2010a. *Neospora caninum* infection in stray and farm dogs: seroepidemiological study and oocyst shedding. *Vet. Parasitol.* 174, 332–335.
- Regidor-Cerrillo, J., Gómez-Bautista, M., Del Pozo, I., Jiménez-Ruiz, E., Aduriz, G., Ortega-Mora, L.M., 2010b. Influence of *Neospora caninum* intra-specific variability in the outcome of infection in a pregnant BALB/c mouse model. *Vet. Res.* 41, 52.
- Regidor-Cerrillo, J., Gomez-Bautista, M., Sodupe, I., Aduriz, G., Álvarez-García, G., Del, P., Ortega-Mora, L.M., 2011. In vitro invasion efficiency and intracellular proliferation rate comprise virulence-related phenotypic traits of *Neospora caninum*. *Vet. Res.* 42, 41.
- Reichel, M.P., Ross, G.P., McAllister, M.M., 2008. Evaluation of an enzyme-linked immunosorbent assay for the serological diagnosis of *Neospora caninum* infection in sheep and determination of the apparent prevalence of infection in New Zealand. *Vet. Parasitol.* 151, 323–326.
- Reichel, M.P., Ellis, J.T., 2009. *Neospora caninum* – How close are we to development of an efficacious vaccine that prevents abortion in cattle? *Int. J. Parasitol.* 39, 1173–1187.

- Reiterová, K., Špilovská, S., Antolová, D., Dubinský, P., 2009. *Neospora caninum*, potential cause of abortions in dairy cows: the current serological follow-up in Slovakia. *Vet. Parasitol.* 159, 1–6.
- Reiterová, K., Špilovská, S., Čobádievová, A., Mucha, R., 2011. First in vitro isolation of *Neospora caninum* from a naturally infected adult dairy cow in Slovakia. *Acta Parasitol.* 56, 111–115.
- Reitt, K., Hilbe, M., Voegtlin, A., Corboz, L., Haessig, M., Pospischil, A., 2007. Aetiology of bovine abortion in Switzerland from 1986 to 1995 – A retrospective study with emphasis on detection of *Neospora caninum* and *Toxoplasma gondii* by PCR. *J. Vet. Med. A.* 54, 15–22.
- Rinaldi, L., Pacelli, F., Iovane, G., Pagnini, U., Veneziano, V., Fusco, G., Cringoli, G., 2007. Survey of *Neospora caninum* and bovine herpes virus 1 coinfection in cattle. *Parasitol. Res.* 100, 359–364.
- Rojo-Montejo, S., Collantes-Fernandez, E., Blanco-Murcia, J., Rodríguez-Bertos, A., Risco-Castillo, V., Ortega-Mora, L.M., 2009a. Experimental infection with a low virulence isolate of *Neospora caninum* at 70 days gestation in cattle did not result in foetopathy. *Vet. Res.* 40, e49.
- Rojo-Montejo, S., Collantes-Fernández, E., Regidor-Cerrillo, J., Álvarez-García, G., Marugán-Hernández, V., Pedraza-Díaz, S., Blanco-Murcia, J., Prenafeta, A., Ortega-Mora, L.M., 2009b. Isolation and characterization of a bovine isolate of *Neospora caninum* with low virulence. *Vet. Parasitol.* 159, 7–16.
- Rojo-Montejo, S., Collantes-Fernández, E., Regidor-Cerrillo, J., Rodríguez-Bertos, A., Prenafeta, A., Gomez-Bautista, M., Ortega-Mora, L.M., 2011. Influence of adjuvant and antigen dose on protection induced by an inactivated whole vaccine against *Neospora caninum* infection in mice. *Vet. Parasitol.* 175, 220–229.
- Romero-Salas, D., García-Vázquez, Z., Montiel-Palacios, F., Montiel-Peña, T., Aguilar-Domínguez, M., Medina-Esparza, L., Cruz-Vázquez, C., 2010. Seroprevalence of *Neospora caninum* antibodies in cattle in Veracruz, Mexico. *J. Anim. Vet. Adv.* 9, 1445–1451.
- Rosbottom, A., Guy, C.S., Gibney, E.H., Smith, R.F., Valarcher, J.F., Taylor, G., Williams, D.J.L., 2007. Peripheral immune responses in pregnant cattle following *Neospora caninum* infection. *Parasite Immunol.* 29, 219–228.
- Rosbottom, A., Gibney, E.H., Guy, C.S., Kipar, A., Smith, R.F., Kaiser, P., Trees, A.J., Williams, D.J.L., 2008. Upregulation of cytokines is detected in the placentas of cattle infected with *Neospora caninum* and is more marked early in gestation when fetal death is observed. *Infect. Immun.* 76, 2352–2361.
- Rosbottom, A., Gibney, H., Kaiser, P., Hartley, C., Smith, R.F., Robinson, R., Kipar, A., Williams, D.J.L., 2011. Up regulation of the maternal immune response in the placenta of cattle naturally infected with *Neospora caninum*. *PLoS ONE* 6, e15799–e15800.
- Rossi, G.F., Cabral, D.D., Ribeiro, D.P., Pajuaba, A.C.A.M., Corrêa, R.R., Moreira, R.Q., Mineo, T.W.P., Mineo, J.R., Silva, D.A.O., 2011. Evaluation of *Toxoplasma gondii* and *Neospora caninum* infections in sheep from Uberlândia, Minas Gerais State Brazil, by different serological methods. *Vet. Parasitol.* 175, 252–259.
- Sadrebazzaz, A., Habibi, G., Haddadzadeh, H., Ashrafi, J., 2007. Evaluation of bovine abortion associated with *Neospora caninum* by different diagnostic techniques in Mashhad, Iran. *Parasitol. Res.* 100, 1257–1260.
- Salaberry, S.R.S., Okuda, L.H., Nassar, A.F.C., de Castro, J.R., Lima-Ribeiro, A.M.C., 2010. Prevalence of *Neospora caninum* antibodies in sheep flocks of Uberlândia county, MG. *Rev. Bras. Parasitol. Vet.* 19, 148–151.
- Salb, A.L., Barkema, H.W., Elkin, B.T., Thompson, R.C.A., Whiteside, D.P., Black, S.R., Dubey, J.P., Kutz, S.J., 2008. Dogs as sources and sentinels of parasites in humans and wildlife, Northern Canada. *Emerg. Infect. Dis.* 14, 60–63.
- Salehi, N., Haddadzadeh, H., Ashrafihelan, J., Shayan, P., Sadrebazzaz, A., 2009. Molecular and pathological study of bovine aborted fetuses and placenta from *Neospora caninum* infected dairy cattle. *Iran. J. Parasitol.* 4, 40–51.
- Sánchez, G.F., Banda, R.V., Sahagun, R.A., Ledesma, M.N., Morales, S.E., 2009. Comparison between immunohistochemistry and two PCR methods for detection of *Neospora caninum* in formalin-fixed and paraffin-embedded brain tissue of bovine fetuses. *Vet. Parasitol.* 164, 328–332.
- Sangster, C., Bryant, B., Campbell-Ward, M., King, J.S., Šlapeta, J., 2010. Neosporosis in an aborted southern white rhinoceros (*Ceratotherium simum simum*) fetus. *J. Zoo Wildlife Med.* 41, 725–728.
- Santolaria, P., López-Gatius, F., Yániz, J., García Ispuerto, I., Nogareda, C., Bech-Sabat, G., Serrano, B., Almería, S., 2009. Early postabortion recovery of *Neospora*-infected lactating dairy cows. *Theriogenology* 72, 798–802.
- Santolaria, P., Almería, S., Martínez-Bello, D., Nogareda, C., Mezo, M., Gonzalez-Warleta, M., Castro-Hermida, J.A., Pabón, M., Yániz, J.L., López-Gatius, F., 2011. Different humoral mechanisms against *Neospora caninum* infection in purebred and crossbreed beef/dairy cattle pregnancies. *Vet. Parasitol.* 178, 70–76.
- Santos, S.L., de Souza Costa, K., Gondim, L.Q., da Silva, M.S.A., Uzêda, R.S., Abe-Sandes, K., Gondim, L.F.P., 2010. Investigation of *Neospora caninum*, *Hammondia* sp., and *Toxoplasma gondii* in tissues from slaughtered beef cattle in Bahia, Brazil. *Parasitol. Res.* 106, 457–461.
- Schares, G., Wilking, H., Bolln, M., Conraths, F.J., Bauer, C., 2009. *Neospora caninum* in dairy herds in Schleswig-Holstein, Germany. *Berl. Münch. Tierärztl. Wochenschr.* 122, 47–50.
- Schwab, A.E., Geary, T.G., Baillargeon, P., Schwab, A.J., Fecteau, G., 2009. Association of BoLA DRB3 and DQA1 alleles with susceptibility to *Neospora caninum* and reproductive outcome in Quebec Holstein cattle. *Vet. Parasitol.* 165, 136–140.
- Scott, H.M., Sorensen, O., Wu, J.T.Y., Chow, E.Y.W., Manninen, K., 2007. Seroprevalence of and agroecological risk factors for *Mycobacterium avium* subspecies *paratuberculosis* and *Neospora caninum* infection among adult beef cattle in cow-calf herds in Alberta, Canada. *Can. Vet. J.* 48, 397–406.
- Segura-Correa, J.C., Domínguez-Díaz, D., Alvalos-Ramírez, R., Argaez-Sosa, J., 2010. Intra-herd correlation coefficients and design effects for bovine viral diarrhoea, infectious bovine rhinotracheitis, leptosporosis and neosporosis in cow-calf system herds in North-eastern Mexico. *Prev. Vet. Med.* 96, 272–275.
- Serrano-Martínez, E., Collantes-Fernández, E., Chávez-Velásquez, A., Rodríguez-Bertos, A., Casas-Astos, E., Risco-Castillo, V., Rosadio-Alcantara, R., Ortega-Mora, L.M., 2007a. Evaluation of *Neospora caninum* and *Toxoplasma gondii* infections in alpaca (*Vicugna pacos*) and llama (*Lama glama*) aborted foetuses from Peru. *Vet. Parasitol.* 150, 39–45.
- Serrano-Martínez, E., Ferre, I., Martínez, A., Osoro, K., Mateos-Sanz, A., del-Pozo, I., Aduriz, G., Tamargo, C., Hidalgo, C.O., Ortega-Mora, L.M., 2007b. Experimental neosporosis in bulls: parasite detection in semen and blood and specific antibody and interferon-gamma responses. *Theriogenology* 67, 1175–1184.
- Serrano, B., Almería, S., García Ispuerto, I., Yániz, J.L., Abdelfattah-Hassan, A., López-Gatius, F., 2011. Peripheral white blood cell counts throughout pregnancy in non-aborting *Neospora caninum*-seronegative and seropositive high-producing dairy cows in a Holstein Friesian herd. *Res. Vet. Sci.* 90, 457–462.
- Shabbir, M.Z., Nazir, M., Lateef, M., Shabbir, M.A.B., Ahmad, A., Rabbani, M., Yaqub, T., Sohail, M.U., Ijaz, M., Maqbool, A. Seroprevalence of *Neospora caninum* and *Brucella abortus* among dairy cattle herds with high abortion rates. *J. Parasitol.*, in press.
- Sharma, S., Bal, M.S., Meenakshi, Kaur, K., Sandhu, K.S., Dubey, J.P., 2008. Seroprevalence of *Neospora caninum* antibodies in dogs in India. *J. Parasitol.* 94, 303–304.
- Silva, M.I.S., Almeida, M.Á.O., Mota, R.A., Pinheiro-Junior, J.W., Rabelo, S.S.A., 2008. Fatores de riscos associados à infecção por *Neospora caninum* em matrizes bovinas leiteiras em Pernambuco. *Ciência Anim. Bras.* 9, 455–461.
- Simek, S., Utuk, A.E., Koroglu, E., Dumanli, N., Risvanli, A., 2008. Seroprevalence of *Neospora caninum* in repeat breeder dairy cows in Turkey. *Arch. Tierz.* 51, 143–148.
- Soares, H.S., Ahid, S.M.M., Bezerra, A.C.D.S., Pena, H.F.J., Dias, R.A., Gennari, S.M., 2009. Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in sheep from Mossoró, Rio Grande do Norte, Brazil. *Vet. Parasitol.* 160, 211–214.
- Soares, R.M., Lopes, E.G., Keid, L.B., Sercundes, M.K., Martins, J., Richtzenhain, L.J., 2011. Identification of *Hammondia heydorni* oocysts by a heminested-PCR (hnPCR-AP10) based on the *H. heydorni* RAPD fragment AP10. *Vet. Parasitol.* 175, 168–172.
- Sobrinho, R., Dubey, J.P., Pabón, M., Linarez, N., Kwok, O.C., Millán, J., Arnal, M.C., Luco, D.F., López-Gatius, F., Thulliez, P., Gortázar, C., Almería, S., 2008. *Neospora caninum* antibodies in wild carnivores from Spain. *Vet. Parasitol.* 155, 190–197.
- Sommanustweechai, A., Vongpakorn, M., Kasantikul, T., Taewnean, J., Siriaronrat, B., Bush, M., Pirarat, N., 2010. Systemic neosporosis in a white rhinoceros. *J. Zoo Wildlife Med.* 41, 165–168.
- Sörgel, S.C., Müller, M., Schares, G., Großmann, E., Neuss, T., Puchta, H., Kreuzer, P., Ewringmann, T., Ehrlein, J., Bogner, K.H., Schmahl, W., 2009. Beteiligung von *Neospora caninum* bei Rinderaborten in Nordbayern. *Tierärztl. Umschau.* 64, 235–243.
- Sotiraki, S., Brozos, C., Samartzi, F., Schares, G., Kioussis, E., Conraths, F.J., 2008. *Neospora caninum* infection in Greek dairy cattle herds detected by two antibody assays in individual milk samples. *Vet. Parasitol.* 152, 79–84.
- Špilovská, S., Reiterová, K., 2008. Seroprevalence of *Neospora caninum* in aborting sheep and goats in the eastern Slovakia. *Folia Vet.* 52, 33–35.

- Špilovská, S., Reiterová, K., Kovacová, D., Bobáková, M., Dubinský, P., 2009. The first finding of *Neospora caninum* and the occurrence of other abortifacient agents in sheep in Slovakia. *Vet. Parasitol.* 164, 320–323.
- Stieve, E., Beckmen, K., Kania, S.A., Widner, A., Patton, S., 2010. *Neospora caninum* and *Toxoplasma gondii* antibody prevalence in Alaska wildlife. *J. Wildlife Dis.* 46, 348–355.
- Strohbusch, M., Müller, N., Hemphill, A., Krebber, R., Greif, G., Gottstein, B., 2009. Toltrazuril treatment of congenitally acquired *Neospora caninum* infection in newborn mice. *Parasitol. Res.* 104, 1335–1343.
- Suteu, O., Titilincu, A., Modry, D., Mihalca, A., Mircean, V., Cozma, V., 2010. First identification of *Neospora caninum* by PCR in aborted bovine foetuses in Romania. *Parasitol. Res.* 106, 719–722.
- Tiwari, A., VanLeeuwen, J.A., Dohoo, I.R., Keefe, G.P., Haddad, J.P., Tremblay, R., Scott, H.M., Whiting, T., 2007. Production effects of pathogens causing bovine leukosis, bovine viral diarrhoea, paratuberculosis, and neosporosis. *J. Dairy Sci.* 90, 659–669.
- Truppel, J.H., Montiani-Ferreira, F., Lange, R.R., Vilani, R.G.O.C., Reifur, L., Boerger, W., da Costa-Ribeiro, M.C.V., Thomaz-Soccol, V., 2010. Detection of *Neospora caninum* DNA in capybaras and phylogenetic analysis. *Parasitol. Int.* 59, 376–379.
- Tschuor, A.C., Kaufmann, T., Strabel, D., Hässig, M., 2010. Abklärung von Aborten und anderen tiergesundheitlichen Problemen beim Rind im Zusammenhang mit der Blauzungenvirus-Impfung 2009. *Schweiz. Arch. Tierheilk.* 152, 501–506.
- Ueno, T.E.H., Gonçalves, V.S.P., Heinemann, M.B., Dilli, T.L.B., Akimoto, B.M., de Souza, S.L.P., Gennari, S.M., Soares, R.M., 2009. Prevalence of *Toxoplasma gondii* and *Neospora caninum* infections in sheep from Federal District, central region of Brazil. *Trop. Anim. Health Prod.* 41, 547–552.
- Uzeda, R.S., Costa, K.S., Santos, S.L., Pinheiro, A.M., de Almeida, M.A.O., McAllister, M.M., Gondim, L.F.P., 2007. Loss of infectivity of *Neospora caninum* oocysts maintained for a prolonged time. *Korean J. Parasitol.* 45, 295–299.
- Valadas, S., Minervino, A.H.H., Lima, V.M.F., Soares, R.M., Ortolani, E.L., Gennari, S.M., 2010a. Occurrence of antibodies anti-*Neospora caninum*, anti-*Toxoplasma gondii*, and anti-*Leishmania chagasi* in serum of dogs from Pará State, Amazon, Brazil. *Parasitol. Res.* 107, 453–457.
- Valadas, S., Gennari, S.M., Yai, L.E.O., Rosypal, A.C., Lindsay, D.S., 2010b. Prevalence of antibodies to *Trypanosoma cruzi*, *Leishmania infantum*, *Encephalitozoon cuniculi*, *Sarcocystis neurona*, and *Neospora caninum* in capybara, *Hydrochoerus hydrochaeris*, from São Paulo State, Brazil. *J. Parasitol.* 96, 521–524.
- Vianna, M.C.B., Sreekumar, C., Miska, K.B., Hill, D.E., Dubey, J.P., 2005. Isolation of *Neospora caninum* from naturally infected white-tailed deer (*Odocoileus virginianus*). *Vet. Parasitol.* 129, 253–257.
- VanLeeuwen, J.A., Haddad, J.P., Dohoo, I.R., Keefe, G.P., Tiwari, A., Scott, H.M., 2010a. Risk factors associated with *Neospora caninum* seropositivity in randomly sampled Canadian dairy cows and herds. *Prev. Vet. Med.* 93, 129–138.
- VanLeeuwen, J.A., Haddad, J.P., Dohoo, I.R., Keefe, G.P., Tiwari, A., Tremblay, R., 2010b. Associations between reproductive performance and seropositivity for bovine leukemia virus, bovine viral diarrhoea virus, *Mycobacterium avium* subspecies *paratuberculosis*, and *Neospora caninum* in Canadian dairy cows. *Prev. Vet. Med.* 94, 54–64.
- VanLeeuwen, J.A., Greenwood, S., Clark, F., Acorn, A., Markham, F., McCarron, J., O'Handley, R., 2011. Monensin use against *Neospora caninum* challenge in dairy cattle. *Vet. Parasitol.* 175, 372–376.
- Vega, O.L., Chávez, V.A., Falcón, P.N., Casas, A.E., 2010. Prevalencia de *Neospora caninum* en perros pastores de una empresa ganadera de la sierra sur del Perú. *Rev. Inv. Vet. Perú.* 21, 80–86.
- Wang, C., Wang, Y., Zou, X., Zhai, Y., Gao, J., Hou, M., Zhu, X.Q., 2010. Seroprevalence of *Neospora caninum* infection in dairy cattle in North-eastern China. *J. Parasitol.* 96, 451–452.
- Wapenaar, W., Barkema, H.W., VanLeeuwen, J.A., McClure, J.T., O'Handley, R.M., Kwok, O.C.H., Thulliez, P., Dubey, J.P., Jenkins, M.C., 2007a. Comparison of serological methods for the diagnosis of *Neospora caninum* infection in cattle. *Vet. Parasitol.* 143, 166–173.
- Wapenaar, W., Barkema, H.W., O'Handley, R.M., Bartels, C.J.M., 2007b. Use of an enzyme-linked immunosorbent assay in bulk milk to estimate the prevalence of *Neospora caninum* on dairy farms in Prince Edward Island, Canada. *Can. Vet. J.* 48, 493–499.
- Wernery, U., Thomas, R., Raghaven, R., Syriac, G., Joseph, S., Georgy, N., 2008. Seroepidemiological studies for the detection of antibodies against 8 infectious diseases in dairy dromedaries of the United Arab Emirates using modern laboratory techniques – part II. *J. Camel Prac. Res.* 15, 139–145.
- West, D.M., Pomroy, W.E., Collett, M.G., Hill, F.I., Ridler, A.L., Kenyon, P.R., Morris, S.T., Pattison, R.S., 2006. A possible role for *Neospora caninum* in ovine abortion in New Zealand. *Small Rumin. Res.* 62, 135–138.
- Weston, J.F., Howe, L., Collett, M.G., Pattison, R.S., Williamson, N.B., West, D.M., Pomroy, W.E., Syed-Hussain, S.S., Morris, S.T., Kenyon, P.R., 2009. Dose-titration challenge of young pregnant sheep with *Neospora caninum* tachyzoites. *Vet. Parasitol.* 164, 183–191.
- Williams, D.J.L., Hartley, C.S., Björkman, C., Trees, A.J., 2009. Endogenous and exogenous transplacental transmission of *Neospora caninum* – how the route of transmission impacts on epidemiology and control of disease. *Parasitology* 136, 1895–1900.
- Woodbine, K.A., Medley, G.F., Moore, S.J., Ramirez-Villaescusa, A., Mason, S., Green, L.E., 2008. A four year longitudinal sero-epidemiology study of *Neospora caninum* in adult cattle from 114 cattle herds in south west England: associations with age, herd and dam-offspring pairs. *BMC Vet. Res.* 4, e35.
- Xia, H.Y., Zhou, D.H., Jia, K., Zeng, X.B., Zhang, D.W., She, L.X., Lin, R.Q., Yuan, Z.G., Li, S.J., 2011. Seroprevalence of *Neospora caninum* infection in dairy cattle of southern China. *J. Parasitol.* 97, 172–173.
- Yai, L.E.O., Ragozo, A.M.A., Cañón-Franco, W.A., Dubey, J.P., Gennari, S.M., 2008. Occurrence of *Neospora caninum* antibodies in capybaras (*Hydrochaeris hydrochaeris*) from São Paulo state, Brazil. *J. Parasitol.* 94, 766.
- Yakhchali, M., Javadi, S., Morshedi, A., 2010. Prevalence of antibodies to *Neospora caninum* in stray dogs of Urmia, Iran. *Parasitol. Res.* 106, 1455–1458.
- Yao, L., Yang, N., Liu, Q., Wang, M., Zhang, W., Qian, W.F., Hu, Y.F., Ding, J., 2009. Detection of *Neospora caninum* in aborted bovine fetuses and dam blood samples by nested PCR and ELISA and seroprevalence in Beijing and Tianjin, China. *Parasitology* 136, 1251–1256.
- Yániz, J.L., López-Gatius, F., García-Ispuerto, I., Bech-Sàbat, G., Serrano, B., Nogareda, C., Sanchez-Nadal, J.A., Almeria, S., Santolaria, P., 2010. Some factors affecting the abortion rate in dairy herds with high incidence of *Neospora*-associated abortions are different in cows and heifers. *Reprod. Domest. Anim.* 45, 699–705.
- Yildiz, K., Kul, O., Babur, C., Kilic, S., Gazyagci, A.N., Celebi, B., Gurcan, I.S., 2009. Seroprevalence of *Neospora caninum* in dairy cattle ranches with high abortion rate: special emphasis to serologic co-existence with *Toxoplasma gondii*, *Brucella abortus* and *Listeria monocytogenes*. *Vet. Parasitol.* 164, 306–310.
- Youssefi, M.R., Arabkhazaeli, F., Hassan, A.T.M., 2009. Seroprevalence of *Neospora caninum* infection in rural and industrial cattle in northern Iran. *Iran. J. Parasitol.* 4, 20–23.
- Yu, X., Chen, N., Hu, D., Zhang, W., Li, X., Wang, B., Kang, L., Li, X., Liu, Q., Tian, K., 2009. Detection of *Neospora caninum* from farm-bred young blue foxes (*Alopex lagopus*) in China. *J. Vet. Med. Sci.* 71, 113–115.
- Zhang, W., Deng, C., Liu, Q., Liu, J., Wang, M., Tian, K.G., Yu, X.L., Hu, D.M., 2007. First identification of *Neospora caninum* infection in aborted bovine foetuses in China. *Vet. Parasitol.* 149, 72–76.